

**DPF2 Polyclonal Antibody**  
Catalog # AP69591**Specification****DPF2 Polyclonal Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">Q92785</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>

**DPF2 Polyclonal Antibody - Additional Information****Gene ID** 5977**Other Names**

DPF2; BAF45D; REQ; UBID4; Zinc finger protein ubi-d4; Apoptosis response zinc finger protein; BRG1-associated factor 45D; BAF45D; D4; zinc and double PHD fingers family 2; Protein requiem

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**DPF2 Polyclonal Antibody - Protein Information****Name** DPF2**Synonyms** BAF45D, REQ, UBID4**Function**

Plays an active role in transcriptional regulation by binding modified histones H3 and H4 (PubMed: [27775714](http://www.uniprot.org/citations/27775714)), PubMed: [28533407](http://www.uniprot.org/citations/28533407)). Is a negative regulator of myeloid differentiation of hematopoietic progenitor cells (PubMed: [28533407](http://www.uniprot.org/citations/28533407)). Might also have a role in the development and maturation of lymphoid cells (By similarity). Involved in the regulation of non-canonical NF-kappa-B pathway (PubMed: [20460684](http://www.uniprot.org/citations/20460684)).

**Cellular Location**

Nucleus. Cytoplasm

## Tissue Location



Ubiquitous.

## DPF2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## DPF2 Polyclonal Antibody - Images



## DPF2 Polyclonal Antibody - Background

Plays an active role in transcriptional regulation by binding modified histones H3 and H4 (PubMed:28533407, PubMed:27775714). Is a negative regulator of myeloid differentiation of hematopoietic progenitor cells (PubMed:28533407). Might also have a role in the development and maturation of lymphoid cells (By similarity). Involved in the regulation of non-canonical NF-kappa-B

pathway (PubMed:20460684).