

**CR2 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6965b**

**Specification**

---

**CR2 Antibody (C-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">P20023</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	986-1014

**CR2 Antibody (C-term) - Additional Information**

**Gene ID** 1380

**Other Names**

Complement receptor type 2, Cr2, Complement C3d receptor, Epstein-Barr virus receptor, EBV receptor, CD21, CR2, C3DR

**Target/Specificity**

This CR2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 986-1014 amino acids from the C-terminal region of human CR2.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CR2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CR2 Antibody (C-term) - Protein Information**

**Name** CR2

**Synonyms** C3DR

**Function** Serves as a receptor for various ligands including complement component CD3d, HNRNPU OR IFNA1 (PubMed:[1849076](#), PubMed:[21527715](#), PubMed:[7753047](#)). When C3d is bound to antigens, attaches to C3d on B- cell surface and thereby facilitates the recognition and uptake of antigens by B-cells (PubMed:[21527715](#)). This interaction enhances B-cell activation and subsequent immune responses. Forms a complex with several partners on the surface of B-cells including CD19, FCRL5 and CD81, to form the B-cell coreceptor complex that plays a crucial role in B-cell activation and signaling (PubMed:[1383329](#), PubMed:[30107486](#)). Induces also specific intracellular signaling separately from the BCR and CD19 by activating the tyrosine kinase SRC, which then phosphorylates nucleolin/NCL and triggers AKT and GSK3 kinase activities in a SYK/CD19-independent manner (PubMed:[12938232](#)). Acts as a ligand for CD23 (FcepsilonRII), a low-affinity receptor for IgE, which is expressed on B-cells and other immune cells, and thus participates in the regulation of IgE production (PubMed:[1386409](#)).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein

#### Tissue Location


Mature B-lymphocytes, T-lymphocytes, pharyngeal epithelial cells, astrocytes and follicular dendritic cells of the spleen

### CR2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### CR2 Antibody (C-term) - Images



Western blot analysis of CR2 Antibody (C-term) (Cat. #AP6965b) in Ramos cell line lysates (35ug/lane). CR2 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human Lymph reacted with CR2 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



CR2 Antibody (C-term) (Cat. #AP6965b) flow cytometry analysis of Ramos cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

**CR2 Antibody (C-term) - Background**

CR2 is a membrane protein, which functions as a receptor for Epstein-Barr virus (EBV) binding on B and T lymphocytes.

**CR2 Antibody (C-term) - References**

Rikova, K., et al., Cell 131 (6), 1190-1203 (2007)