

**eIF3 $\alpha$  Polyclonal Antibody**  
Catalog # AP69695**Specification****eIF3 $\alpha$  Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O75822</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**eIF3 $\alpha$  Polyclonal Antibody - Additional Information**

Gene ID 8669

**Other Names**

EIF3J; EIF3S1; Eukaryotic translation initiation factor 3 subunit J; eIF3j; Eukaryotic translation initiation factor 3 subunit 1; eIF-3-alpha; eIF3 p35

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**eIF3 $\alpha$  Polyclonal Antibody - Protein Information**

Name EIF3J {ECO:0000255|HAMAP-Rule:MF\_03009}

**Function**

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:<a href="http://www.uniprot.org/citations/25849773" target="\_blank">25849773</a>, PubMed:<a href="http://www.uniprot.org/citations/27462815" target="\_blank">27462815</a>). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:<a href="http://www.uniprot.org/citations/25849773" target="\_blank">25849773</a>).


**Cellular Location**

Cytoplasm {ECO:0000255|HAMAP-Rule:MF\_03009}.

**eIF3 $\alpha$  Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**eIF3 $\alpha$  Polyclonal Antibody - Images****eIF3 $\alpha$  Polyclonal Antibody - Background**

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA<sub>i</sub> and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination

ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773).