

EIF5B Polyclonal Antibody
Catalog # AP69704**Specification**

EIF5B Polyclonal Antibody - Product Information

Application	WB
Primary Accession	O60841
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

EIF5B Polyclonal Antibody - Additional Information**Gene ID** 9669**Other Names**

EIF5B; IF2; KIAA0741; Eukaryotic translation initiation factor 5B; eIF-5B; Translation initiation factor IF-2

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

EIF5B Polyclonal Antibody - Protein Information**Name** EIF5B**Synonyms** IF2, KIAA0741**Function**

Plays a role in translation initiation (PubMed:10659855, PubMed:35732735). Ribosome-dependent GTPase that promotes the joining of the 60S ribosomal subunit to the pre-initiation complex to form the 80S initiation complex with the initiator methionine-tRNA in the P-site base paired to the start codon (PubMed:10659855, PubMed:35732735). Together with eIF1A (EIF1AX), actively orients the initiator methionine-tRNA in a conformation that allows 60S ribosomal subunit joining to form the 80S initiation complex (PubMed:12569173, PubMed:35732735). Is released after formation of the 80S initiation complex (PubMed:12569173).

[35732735](http://www.uniprot.org/citations/35732735)). Its GTPase activity is not essential for ribosomal subunits joining, but GTP hydrolysis is needed for eIF1A (EIF1AX) ejection quickly followed by EIF5B release to form elongation-competent ribosomes (PubMed: [10659855](http://www.uniprot.org/citations/10659855), PubMed: [35732735](http://www.uniprot.org/citations/35732735)). In contrast to its procaryotic homolog, does not promote recruitment of Met-rRNA to the small ribosomal subunit (PubMed: [10659855](http://www.uniprot.org/citations/10659855)).

Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:Q05D44}.

eIF5B Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

eIF5B Polyclonal Antibody - Images



eIF5B Polyclonal Antibody - Background

Plays a role in translation initiation. Translational GTPase that catalyzes the joining of the 40S and 60S subunits to form the 80S initiation complex with the initiator methionine-tRNA in the P-site base paired to the start codon. GTP binding and hydrolysis induces conformational changes in the enzyme that renders it active for productive interactions with the ribosome. The release of the

enzyme after formation of the initiation complex is a prerequisite to form elongation-competent ribosomes.