

**HSF-27 Polyclonal Antibody**  
Catalog # AP70421**Specification****HSF-27 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O96JB5</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**HSF-27 Polyclonal Antibody - Additional Information**

Gene ID 80279

**Other Names**

CDK5RAP3; IC53; MSTP016; OK/SW-cl.114; PP1553; CDK5 regulatory subunit-associated protein 3; CDK5 activator-binding protein C53; Protein HSF-27

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**HSF-27 Polyclonal Antibody - Protein Information**Name CDK5RAP3 ([HGNC:18673](#))**Function**

Substrate adapter for ufmylation, the covalent attachment of the ubiquitin-like modifier UFM1 to substrate proteins, in response to endoplasmic reticulum stress (PubMed: [23152784](http://www.uniprot.org/citations/23152784), PubMed: [30635284](http://www.uniprot.org/citations/30635284)). Negatively regulates NF-kappa-B-mediated gene transcription through the control of RELA phosphorylation (PubMed: [17785205](http://www.uniprot.org/citations/17785205), PubMed: [20228063](http://www.uniprot.org/citations/20228063)). Probable tumor suppressor initially identified as a CDK5R1 interactor controlling cell proliferation (PubMed: [12054757](http://www.uniprot.org/citations/12054757), PubMed: [12737517](http://www.uniprot.org/citations/12737517)). Also regulates mitotic G2/M transition checkpoint and mitotic G2 DNA damage checkpoint (PubMed: [15790566](http://www.uniprot.org/citations/15790566), PubMed: [19223857](http://www.uniprot.org/citations/19223857)). Through its interaction with CDKN2A/ARF and MDM2 may induce MDM2-dependent p53/TP53

ubiquitination, stabilization and activation in the nucleus, thereby promoting G1 cell cycle arrest and inhibition of cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/16173922" target="\_blank">16173922</a>). May also play a role in the rupture of the nuclear envelope during apoptosis (PubMed:<a href="http://www.uniprot.org/citations/23478299" target="\_blank">23478299</a>). May regulate MAPK14 activity by regulating its dephosphorylation by PPM1D/WIP1 (PubMed:<a href="http://www.uniprot.org/citations/21283629" target="\_blank">21283629</a>). Required for liver development (By similarity).

#### Cellular Location

Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome  
Note=Colocalizes and associates with microtubules

#### Tissue Location


Ubiquitously expressed (PubMed:10721722, PubMed:12054757). Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Isoform 3 is expressed in kidney, liver, skeletal muscle and placenta (PubMed:12737517)




### HSF-27 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### HSF-27 Polyclonal Antibody - Images





### HSF-27 Polyclonal Antibody - Background

Probable tumor suppressor initially identified as a CDK5R1 interactor controlling cell proliferation (PubMed:12054757, PubMed:12737517). Negatively regulates NF-kappa-B-mediated gene transcription through the control of RELA phosphorylation (PubMed:17785205, PubMed:20228063). Also regulates mitotic G2/M transition checkpoint and mitotic G2 DNA damage checkpoint (PubMed:15790566, PubMed:19223857). Through its interaction with CDKN2A/ARF and MDM2 may induce MDM2-dependent p53/TP53 ubiquitination, stabilization and activation in the nucleus, thereby promoting G1 cell cycle arrest and inhibition of cell proliferation (PubMed:16173922). May play a role in the unfolded protein response, mediating the ufmylation of multiple proteins in response to endoplasmic reticulum stress (PubMed:23152784). May also play a role in the rupture of the nuclear envelope during apoptosis (PubMed:23478299). May regulate MAPK14 activity by regulating its dephosphorylation by PPM1D/WIP1 (PubMed:21283629).