

IL-4R α Polyclonal Antibody
Catalog # AP70520**Specification**

IL-4R α Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P24394
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

IL-4R α Polyclonal Antibody - Additional Information**Gene ID** 3566**Other Names**

IL4R; IL4RA; 582J2.1; Interleukin-4 receptor subunit alpha; IL-4 receptor subunit alpha; IL-4R subunit alpha; IL-4R-alpha; IL-4RA; CD antigen CD124

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

IL-4R α Polyclonal Antibody - Protein Information**Name** IL4R**Synonyms** IL4RA**Function**

Receptor for both interleukin 4 and interleukin 13. Couples to the JAK1/2/3-STAT6 pathway. The IL4 response is involved in promoting Th2 differentiation. The IL4/IL13 responses are involved in regulating IgE production and, chemokine and mucus production at sites of allergic inflammation. In certain cell types, can signal through activation of insulin receptor substrates, IRS1/IRS2.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location



Isoform 1 and isoform 2 are highly expressed in activated T-cells

IL-4R α Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL-4R α Polyclonal Antibody - Images



IL-4R α Polyclonal Antibody - Background

Receptor for both interleukin 4 and interleukin 13. Couples to the JAK1/2/3-STAT6 pathway. The IL4

response is involved in promoting Th2 differentiation. The IL4/IL13 responses are involved in regulating IgE production and, chemokine and mucus production at sites of allergic inflammation. In certain cell types, can signal through activation of insulin receptor substrates, IRS1/IRS2.