

LT- β Polyclonal Antibody
Catalog # AP70784**Specification**

LT- β Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q06643
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

LT- β Polyclonal Antibody - Additional Information**Gene ID** 4050**Other Names**

LTB; TNFC; TNFSF3; Lymphotoxin-beta; LT-beta; Tumor necrosis factor C; TNF-C; Tumor necrosis factor ligand superfamily member 3

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

LT- β Polyclonal Antibody - Protein Information**Name** LTB**Synonyms** TNFC, TNFSF3**Function**

Cytokine that binds to LTBR/TNFRSF3. May play a specific role in immune response regulation. Provides the membrane anchor for the attachment of the heterotrimeric complex to the cell surface. Isoform 2 is probably non-functional.

Cellular Location

Membrane; Single-pass type II membrane protein

Tissue Location



Spleen and thymus.

LT- β Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LT- β Polyclonal Antibody - Images



LT- β Polyclonal Antibody - Background

Cytokine that binds to LTBR/TNFRSF3. May play a specific role in immune response regulation.

Provides the membrane anchor for the attachment of the heterotrimeric complex to the cell surface. Isoform 2 is probably non-functional.