

**Midline-1 Polyclonal Antibody**  
Catalog # AP70943

**Specification**

---

**Midline-1 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O15344</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**Midline-1 Polyclonal Antibody - Additional Information**

**Gene ID** 4281

**Other Names**

MID1; FXY; RNF59; TRIM18; XPRF; Midline-1; Midin; Midline 1 RING finger protein; Putative transcription factor XPRF; RING finger protein 59; Tripartite motif-containing protein 18

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Midline-1 Polyclonal Antibody - Protein Information**

**Name** MID1

**Synonyms** FXY, RNF59, TRIM18, XPRF

**Function**

Has E3 ubiquitin ligase activity towards IGBP1, promoting its monoubiquitination, which results in deprotection of the catalytic subunit of protein phosphatase PP2A, and its subsequent degradation by polyubiquitination.

**Cellular Location**

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, spindle. Note=Microtubule-associated. It is associated with microtubules throughout the cell cycle, co-localizing with cytoplasmic fibers in interphase and with the mitotic spindle and midbodies during mitosis and cytokinesis

**Tissue Location**

In the fetus, highest expression found in kidney, followed by brain and lung. Expressed at low



levels in fetal liver. In the adult, most abundant in heart, placenta and brain

### Midline-1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Midline-1 Polyclonal Antibody - Images



**Midline-1 Polyclonal Antibody - Background**

Has E3 ubiquitin ligase activity towards IGBP1, promoting its monoubiquitination, which results in deprotection of the catalytic subunit of protein phosphatase PP2A, and its subsequent degradation by polyubiquitination.