

NCoA-7 Polyclonal Antibody
Catalog # AP71181**Specification**

NCoA-7 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q8NI08
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

NCoA-7 Polyclonal Antibody - Additional Information**Gene ID** 135112**Other Names**

NCOA7; ERAP140; ESNA1; Nbla00052; Nbla10993; Nuclear receptor coactivator 7; 140 kDa estrogen receptor-associated protein; Estrogen nuclear receptor coactivator 1

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

NCoA-7 Polyclonal Antibody - Protein Information**Name** NCOA7**Synonyms** ERAP140, ESNA1**Function**

Enhances the transcriptional activities of several nuclear receptors. Involved in the coactivation of different nuclear receptors, such as ESR1, THRB, PPARG and RARA.

Cellular Location

Nucleus.

Tissue Location


Highly expressed in brain. Weakly expressed in mammary gland, ovary, uterus, prostate, stomach, bladder, spinal cord and pancreas. Expressed in cancer cell line

NCoA-7 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCoA-7 Polyclonal Antibody - Images



NCoA-7 Polyclonal Antibody - Background

Enhances the transcriptional activities of several nuclear receptors. Involved in the coactivation of different nuclear receptors, such as ESR1, THRB, PPARG and RARA.