

ND5 Polyclonal Antibody
Catalog # AP71184**Specification**

ND5 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P03915
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

ND5 Polyclonal Antibody - Additional Information**Gene ID** 4540**Other Names**

MT-ND5; MTND5; NADH5; ND5; NADH-ubiquinone oxidoreductase chain 5; NADH dehydrogenase subunit 5

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

ND5 Polyclonal Antibody - Protein Information**Name** MT-ND5**Synonyms** MTND5, NADH5, ND5**Function**

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (PubMed:15250827). Essential for the catalytic activity and assembly of complex I (PubMed:15250827).

Cellular Location


Mitochondrion inner membrane {ECO:0000250|UniProtKB:P03920}; Multi-pass membrane protein

ND5 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ND5 Polyclonal Antibody - Images



ND5 Polyclonal Antibody - Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).