

NDUFV2 Polyclonal Antibody
Catalog # AP71205**Specification****NDUFV2 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	P19404
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

NDUFV2 Polyclonal Antibody - Additional Information**Gene ID** 4729**Other Names**

NDUFV2; NADH dehydrogenase [ubiquinone] flavoprotein 2; mitochondrial; NADH-ubiquinone oxidoreductase 24 kDa subunit

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

NDUFV2 Polyclonal Antibody - Protein Information**Name** NDUFV2 ([HGNC:7717](#))**Function**

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (Probable). Parts of the peripheral arm of the enzyme, where the electrons from NADH are accepted by flavin mononucleotide (FMN) and then passed along a chain of iron-sulfur clusters by electron tunnelling to the final acceptor ubiquinone (Probable). Contains one iron-sulfur cluster (Probable).

Cellular Location



Mitochondrion inner membrane {ECO:0000250|UniProtKB:P04394}; Peripheral membrane protein {ECO:0000250|UniProtKB:P04394}; Matrix side {ECO:0000250|UniProtKB:P04394}


NDUFV2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NDUFV2 Polyclonal Antibody - Images





NDUFV2 Polyclonal Antibody - Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).