

Neuropsin Polyclonal Antibody
Catalog # AP71259**Specification****Neuropsin Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	Q6U736
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

Neuropsin Polyclonal Antibody - Additional Information**Gene ID** 221391**Other Names**

OPN5; GPR136; PGR12; TMEM13; Opsin-5; G-protein coupled receptor 136; G-protein coupled receptor PGR12; Neuropsin; Transmembrane protein 13

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Neuropsin Polyclonal Antibody - Protein Information**Name** OPN5**Synonyms** GPR136, PGR12, TMEM13**Function**

G-protein coupled receptor which selectively activates G(i) type G proteins via ultraviolet A (UVA) light-mediated activation in the retina (By similarity). Preferentially binds the chromophore 11-cis retinal and is a bistable protein that displays emission peaks at 380 nm (UVA light) and 470 nm (blue light) (PubMed:22043319). Required for the light-response in the inner plexiform layer, and contributes to the regulation of the light-response in the nerve fiber layer, via phosphorylated DAT/SLC6A3 dopamine uptake (By similarity). Involved in local corneal and retinal circadian rhythm photoentrainment via modulation of the UVA light-induced phase-shift of the retina clock (By similarity). Acts as a circadian photoreceptor in the outer ear, via modulation of circadian clock-gene expression in response to violet light during the light-to-dark transition phase and night phase of the circadian cycle (By similarity). Required in the retina to negatively regulate hyaloid vessel regression during postnatal development via light-dependent OPN5-SLC32A1-DRD2-VEGFR2

signaling (By similarity). Involved in the light-dependent regulation of retina and vitreous compartment dopamine levels (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

Detected in brain and retina and cell lines derived from neural retina.

Neuropsin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Neuropsin Polyclonal Antibody - Images



