

## PDXK Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7167a

### Specification

## PDXK Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	<u>000764</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35102
Antigen Region	242-272

## PDXK Antibody (C-term) - Additional Information

Gene ID 8566

Other Names Pyridoxal kinase, Pyridoxine kinase, PDXK, C21orf124, C21orf97, PKH, PNK

Target/Specificity

This PDXK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 242-272 amino acids from the C-terminal region of human PDXK.

**Dilution** WB~~1:1000 IHC-P~~1:50~100

### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

PDXK Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## PDXK Antibody (C-term) - Protein Information

Name PDXK (<u>HGNC:8819</u>)

**Function** Catalyzes the phosphorylation of the dietary vitamin B6 vitamers pyridoxal (PL), pyridoxine (PN) and pyridoxamine (PM) to form pyridoxal 5'-phosphate (PLP), pyridoxine



5'-phosphate (PNP) and pyridoxamine 5'-phosphate (PMP), respectively (Probable) (PubMed:<u>10987144</u>, PubMed:<u>17766369</u>, PubMed:<u>19351586</u>, PubMed:<u>31187503</u>, PubMed:<u>9099727</u>). PLP is the active form of vitamin B6, and acts as a cofactor for over 140 different enzymatic reactions.


**Cellular Location** Cytoplasm, cytosol.

**Tissue Location** Ubiquitous (PubMed:31187503, PubMed:9099727). Highly expressed in testis (PubMed:9099727)

# PDXK Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- PDXK Antibody (C-term) Images



Western blot analysis of anti-PDXK (Cat. #AP7167a) in HepG2 cell line lysate (35ug/lane).PDXK(arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

# PDXK Antibody (C-term) - Background

Pyridoxal kinase (PDXK) converts vitamin B6 to pyridoxal-5-phosphate (PLP), an essential cofactor in the intermediate metabolism of amino acids and neurotransmitters. The PDXK gene encodes a 312-amino acid polypeptide, and expression of the cDNA reveals pyridoxal kinase activity. Northern blot analysis revealed that a major 1.5-kb PDXK transcript is expressed in all tissues tested. The expression of PDXK shows circadian oscillations. The expression of Pdxk in mouse liver and brain is regulated by the 3 PAR bZIP transcription factors, Dbp, Hlf, and Tef, which also show circadian oscillations in expression. Mice devoid of all 3 transcription factors show decreased levels of brain PLP, serotonin, and dopamine, and are highly susceptible to frequently lethal generalized spontaneous and audiogenic epilepsies.

## PDXK Antibody (C-term) - References

Shin, J.H., et al., Neurochem. Int. 45(1):73-79 (2004). Lee, H.S., et al., Mol. Cells 10(4):452-459 (2000). Laine-Cessac, P., et al., Biochem. Pharmacol. 54(8):863-870 (1997). Hanna, M.C., et al., J. Biol. Chem. 272(16):10756-10760 (1997). Zhang, Z., et al., J. Nutr. 10(1):53-59 (1993). **PDXK Antibody (C-term) - Citations** 

• <u>Metabolic features of cancer cells impact immunosurveillance</u>