

PBK Polyclonal Antibody
Catalog # AP71794**Specification**

PBK Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q96KB5
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal

PBK Polyclonal Antibody - Additional Information**Gene ID** 55872**Other Names**

PBK; TOPK; Lymphokine-activated killer T-cell-originated protein kinase; Cancer/testis antigen 84; CT84; MAPKK-like protein kinase; Nori-3; PDZ-binding kinase; Spermatogenesis-related protein kinase; SPK; T-LAK cell-originated protein kinas

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

PBK Polyclonal Antibody - Protein Information**Name** PBK**Synonyms** TOPK**Function**

Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin- induced DNA damage.

Tissue Location


Expressed in the testis and placenta. In the testis, restrictedly expressed in outer cell layer of seminiferous tubules.

PBK Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PBK Polyclonal Antibody - Images



PBK Polyclonal Antibody - Background

Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin-induced DNA damage.