

**CAMK2G (CAMK2 gamma) Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7208a**

**Specification**

---

**CAMK2G (CAMK2 gamma) Antibody (C-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">O13555</a>
Other Accession	<a href="#">O15378</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	309-338

**CAMK2G (CAMK2 gamma) Antibody (C-term) - Additional Information**

**Gene ID** 818

**Other Names**

Calcium/calmodulin-dependent protein kinase type II subunit gamma, CaM kinase II subunit gamma, CaMK-II subunit gamma, CAMK2G, CAMK, CAMK-II, CAMKG

**Target/Specificity**

This CAMK2G (CAMK2 gamma) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 309-338 amino acids from the C-terminal region of human CAMK2G (CAMK2 gamma).

**Dilution**

WB~~1:2000  
IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CAMK2G (CAMK2 gamma) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CAMK2G (CAMK2 gamma) Antibody (C-term) - Protein Information**

**Name** CAMK2G

**Synonyms** CAMK, CAMK-II, CAMKG

**Function** Calcium/calmodulin-dependent protein kinase that functions autonomously after Ca(2+)/calmodulin-binding and autophosphorylation, and is involved in sarcoplasmic reticulum Ca(2+) transport in skeletal muscle and may function in dendritic spine and synapse formation and neuronal plasticity (PubMed:[16690701](#)). In slow-twitch muscles, is involved in regulation of sarcoplasmic reticulum (SR) Ca(2+) transport and in fast-twitch muscle participates in the control of Ca(2+) release from the SR through phosphorylation of the ryanodine receptor-coupling factor triadin (PubMed:[16690701](#)). In the central nervous system, it is involved in the regulation of neurite formation and arborization (PubMed:[30184290](#)). It may participate in the promotion of dendritic spine and synapse formation and maintenance of synaptic plasticity which enables long-term potentiation (LTP) and hippocampus-dependent learning. In response to interferon-gamma (IFN-gamma) stimulation, catalyzes phosphorylation of STAT1, stimulating the JAK-STAT signaling pathway (By similarity).

**Cellular Location**

Sarcoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side


**Tissue Location**

Expressed in skeletal muscle.


**CAMK2G (CAMK2 gamma) Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CAMK2G (CAMK2 gamma) Antibody (C-term) - Images**


The anti-CAMK2 gamma C-term Pab (Cat. #AP7208a) is used in Western blot to detect CAMK2 gamma in mouse heart tissue lysate.




All lanes : Anti-CAMK2G (CAMK2 gamma) Antibody (C-term) at 1:500-1000 dilution Lane 1: Jurkat whole cell lysate Lane 2: U-251 MG whole cell lysate Lane 3: Human brain tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-CAMK2G (CAMK2 gamma) Antibody (C-term) at 1:500 dilution + T47D whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-CAMK2G (CAMK2 gamma) Antibody (C-term) at 1:2000 dilution + human liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 63kDa Blocking/Dilution buffer: 5% NFD/MTBST.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



All lanes : Anti-CAMK2G (CAMK2 gamma) Antibody (C-term) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),

Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

### **CAMK2G (CAMK2 gamma) Antibody (C-term) - Background**

CaM-kinase II (CAMK2) is a prominent Ser/Thr protein kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Likely autophosphorylation of Thr-286 allows the kinase to switch from a calmodulin-dependent to a calmodulin-independent state. CAMK2 is composed of four different chains: alpha, beta, gamma, and delta. The different isoforms assemble into homo- or heteromultimeric holoenzymes composed of 8 to 12 subunits.

### **CAMK2G (CAMK2 gamma) Antibody (C-term) - References**

Breen, M.A., et al., Biochem. Biophys. Res. Commun. 236(2):473-478 (1997). Tombes, R.M., et al., Biochim. Biophys. Acta 1355(3):281-292 (1997).

### **CAMK2G (CAMK2 gamma) Antibody (C-term) - Citations**

- [Matrine inhibits the growth of natural killer/T-cell lymphoma cells by modulating CaMKII \$\gamma\$ -c-Myc signaling pathway](#)
- [Activation of CaMKII \$\gamma\$  potentiates T-cell acute lymphoblastic leukemia leukemogenesis via phosphorylating FOXO3a.](#)