

**SPHK2 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7238B**

**Specification**

---

**SPHK2 Antibody (C-term) - Product Information**

Application	WB, IHC-P-Leica,E
Primary Accession	<a href="#">O9NRA0</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	590-620

**SPHK2 Antibody (C-term) - Additional Information**

**Gene ID** 56848

**Other Names**

Sphingosine kinase 2, SK 2, SPK 2, SPHK2

**Target/Specificity**

This SPHK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 590-620 amino acids from the C-terminal region of human SPHK2.

**Dilution**

WB~~1:1000  
IHC-P-Leica~~1:250

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SPHK2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**SPHK2 Antibody (C-term) - Protein Information**

**Name** SPHK2 ([HGNC:18859](#))

**Synonyms** SK2

**Function** Catalyzes the phosphorylation of sphingosine to form sphingosine-1-phosphate (SPP), a

lipid mediator with both intra- and extracellular functions. Also acts on D-erythro-dihydrosphingosine, D- erythro-sphingosine and L-threo-dihydrosphingosine. Binds phosphoinositides (PubMed:[12954646](#), PubMed:[19168031](#)). In contrast to prosurvival SPHK1, has a positive effect on intracellular ceramide levels, inhibits cells growth and enhances apoptosis (PubMed:[16118219](#)). In mitochondria, is important for cytochrome-c oxidase assembly and mitochondrial respiration. The SPP produced in mitochondria binds PHB2 and modulates the regulation via PHB2 of complex IV assembly and respiration (PubMed:[20959514](#)). In nucleus, plays a role in epigenetic regulation of gene expression. Interacts with HDAC1 and HDAC2 and, through SPP production, inhibits their enzymatic activity, preventing the removal of acetyl groups from lysine residues with histones. Up- regulates acetylation of histone H3-K9, histone H4-K5 and histone H2B- K12 (PubMed:[19729656](#)). In nucleus, may have an inhibitory effect on DNA synthesis and cell cycle (PubMed:[12954646](#), PubMed:[16103110](#)). In mast cells, is the main regulator of SPP production which mediates calcium influx, NF-kappa-B activation, cytokine production, such as TNF and IL6, and degranulation of mast cells (By similarity). In dopaminergic neurons, is involved in promoting mitochondrial functions regulating ATP and ROS levels (By similarity). Also involved in the regulation of glucose and lipid metabolism (By similarity).

#### Cellular Location

Cytoplasm. Nucleus. Endoplasmic reticulum {ECO:0000250|UniProtKB:Q9JIA7}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q9JIA7}. Note=In nucleus, located in nucleosomes where it associates with core histone proteins such as histone 3 (PubMed:[19729656](#)). In brains of patients with Alzheimer's disease, may be preferentially localized in the nucleus. Cytosolic expression decrease correlates with the density of amyloid deposits (PubMed:[29615132](#)). In apoptotic cells, colocalizes with CASP1 in cell membrane where is cleaved and released from cells in an active form (PubMed:[20197547](#)).

#### Tissue Location

Mainly expressed in adult kidney, liver, and brain (PubMed:[10751414](#)). Expressed in cerebral cortex and hippocampus (at protein level) (PubMed:[29615132](#)). Isoform 1 is the predominant form expressed in most tissues (PubMed:[16103110](#))


#### SPHK2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### SPHK2 Antibody (C-term) - Images







All lanes : Anti-hSPHK2-A605 at 1:1000 dilution Lane 1: HepG2 whole cell lysate Lane 2: A549 whole cell lysate Lane 3: 293 whole cell lysate Lane 4: U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 69 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



SPHK2 Antibody (A605) (Cat. #AP7238b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the SPHK2 antibody detected the SPHK2 protein (arrow).



Immunohistochemical analysis of AP7238B on paraffin-embedded Human brain tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:250) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.



Immunohistochemical analysis of AP7238B on paraffin-embedded Human kidney tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:250) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.

### **SPHK2 Antibody (C-term) - Background**

Sphingosine Kinase (SphK) catalyzes the phosphorylation of the lipid sphingosine, creating the bioactive lipid sphingosine-1-phosphate (S1P). S1P subsequently signals through cell surface G protein-coupled receptors, as well as intracellularly, to modulate cell proliferation, survival, motility and differentiation. SphK is an important signaling enzyme which is activated by diverse agents, including growth factors that signal through receptor tyrosine kinases, agents activating G protein-coupled receptors, and immunoglobulin receptors. Two SphK isotypes, SphK-1 and SphK-2, have been cloned, and both isotypes are ubiquitously expressed. SphK-1 has been shown to mediate cell growth, prevention of apoptosis, and cellular transformation, and is upregulated in a

variety of human tumors. In contrast, SphK-2 increases apoptosis, and may be responsible for phosphorylating and activating the immunosuppressive drug FTY720.

#### **SPHK2 Antibody (C-term) - References**

Shu, X., et al., Mol. Cell. Biol. 22(22):7758-7768 (2002).  
Xia, P., et al., J. Biol. Chem. 277(10):7996-8003 (2002).  
Liu, H., et al., J. Biol. Chem. 275(26):19513-19520 (2000).  
Shutler, G., et al., Genomics 34(3):334-339 (1996).

#### **SPHK2 Antibody (C-term) - Citations**

- [Targeting sphingosine kinase 2 suppresses cell growth and synergizes with BCL2/BCL-XL inhibitors through NOXA-mediated MCL1 degradation in cholangiocarcinoma.](#)
- [Down-regulation of Sphk2 suppresses bladder cancer progression.](#)
- [Sphingosine 1-phosphate modulates spinal nociceptive processing.](#)