

SENP2 Polyclonal Antibody
Catalog # AP72420**Specification****SENP2 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	Q9HC62
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

SENP2 Polyclonal Antibody - Additional Information

Gene ID 59343

Other Names

SENP2; KIAA1331; Sentrin-specific protease 2; Axam2; SMT3-specific isopeptidase 2; Smt3ip2; Sentrin/SUMO-specific protease SENP2

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

SENP2 Polyclonal Antibody - Protein Information**Name** SENP2 {ECO:0000303|PubMed:10718198, ECO:0000312|HGNC:HGNC:23116}**Function**

Protease that catalyzes two essential functions in the SUMO pathway (PubMed:[11896061](http://www.uniprot.org/citations/11896061), PubMed:[12192048](http://www.uniprot.org/citations/12192048), PubMed:[15296745](http://www.uniprot.org/citations/15296745), PubMed:[20194620](http://www.uniprot.org/citations/20194620), PubMed:[21965678](http://www.uniprot.org/citations/21965678)). The first is the hydrolysis of an alpha-linked peptide bond at the C-terminal end of the small ubiquitin-like modifier (SUMO) propeptides, SUMO1, SUMO2 and SUMO3 leading to the mature form of the proteins (PubMed:[15296745](http://www.uniprot.org/citations/15296745)). The second is the deconjugation of SUMO1, SUMO2 and SUMO3 from targeted proteins, by cleaving an epsilon-linked peptide bond between the C-terminal glycine of the mature SUMO and the lysine epsilon-amino group of the target protein (PubMed:[15296745](http://www.uniprot.org/citations/15296745), PubMed:[20194620](http://www.uniprot.org/citations/20194620), PubMed:[20194620](http://www.uniprot.org/citations/20194620), PubMed:[20194620](http://www.uniprot.org/citations/20194620)).

<http://www.uniprot.org/citations/21965678> target="_blank">21965678). May down-regulate CTNNB1 levels and thereby modulate the Wnt pathway (By similarity). Deconjugates SUMO2 from MTA1 (PubMed:21965678). Plays a dynamic role in adipogenesis by desumoylating and promoting the stabilization of CEBPB (PubMed:20194620). Acts as a regulator of the cGAS-STING pathway by catalyzing desumoylation of CGAS and STING1 during the late phase of viral infection (By similarity).

Cellular Location


Nucleus, nuclear pore complex. Nucleus membrane; Peripheral membrane protein; Nucleoplasmic side. Cytoplasm Note=Shuttles between cytoplasm and nucleus

SEN2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SEN2 Polyclonal Antibody - Images



SEN2 Polyclonal Antibody - Background

Protease that catalyzes two essential functions in the SUMO pathway. The first is the hydrolysis of an alpha-linked peptide bond at the C-terminal end of the small ubiquitin-like modifier (SUMO) propeptides, SUMO1, SUMO2 and SUMO3 leading to the mature form of the proteins. The second is the deconjugation of SUMO1, SUMO2 and SUMO3 from targeted proteins, by cleaving an epsilon-linked peptide bond between the C-terminal glycine of the mature SUMO and the lysine

epsilon-amino group of the target protein. May down-regulate CTNNB1 levels and thereby modulate the Wnt pathway. Deconjugates SUMO2 from MTA1. Plays a dynamic role in adipogenesis by desumoylating and promoting the stabilization of CEBPB (PubMed:20194620).