

**SH-PTP1 Polyclonal Antibody**  
Catalog # AP72474**Specification**

---

**SH-PTP1 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P29350</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**SH-PTP1 Polyclonal Antibody - Additional Information****Gene ID** 5777**Other Names**

PTPN6; HCP; PTP1C; Tyrosine-protein phosphatase non-receptor type 6; Hematopoietic cell protein-tyrosine phosphatase; Protein-tyrosine phosphatase 1C; PTP-1C; Protein-tyrosine phosphatase SHP-1; SH-PTP1

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**SH-PTP1 Polyclonal Antibody - Protein Information****Name** PTPN6**Synonyms** HCP, PTP1C**Function**

Modulates signaling by tyrosine phosphorylated cell surface receptors such as KIT and the EGF receptor/EGFR. The SH2 regions may interact with other cellular components to modulate its own phosphatase activity against interacting substrates. Together with MTUS1, induces UBE2V2 expression upon angiotensin II stimulation. Plays a key role in hematopoiesis.

**Cellular Location**

Cytoplasm. Nucleus. Note=In neurons, translocates into the nucleus after treatment with angiotensin II (By similarity) Shuttles between the cytoplasm and nucleus via its association with PDPK1.

**Tissue Location**



Isoform 1 is expressed in hematopoietic cells. Isoform 2 is expressed in non-hematopoietic cells

### SH-PTP1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### SH-PTP1 Polyclonal Antibody - Images



### SH-PTP1 Polyclonal Antibody - Background

Modulates signaling by tyrosine phosphorylated cell surface receptors such as KIT and the EGF receptor/EGFR. The SH2 regions may interact with other cellular components to modulate its own phosphatase activity against interacting substrates. Together with MTUS1, induces UBE2V2 expression upon angiotensin II stimulation. Plays a key role in hematopoiesis.