

SH-PTP2 Polyclonal Antibody
Catalog # AP72477**Specification****SH-PTP2 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	Q06124
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

SH-PTP2 Polyclonal Antibody - Additional Information

Gene ID 5781

Other Names

PTPN11; PTP2C; SHPTP2; Tyrosine-protein phosphatase non-receptor type 11; Protein-tyrosine phosphatase 1D; PTP-1D; Protein-tyrosine phosphatase 2C; PTP-2C; SH-PTP2; SHP-2; Shp2; SH-PTP3

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

SH-PTP2 Polyclonal Antibody - Protein Information

Name PTPN11

Synonyms PTP2C, SHPTP2

Function

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus (PubMed:[10655584](http://www.uniprot.org/citations/10655584)), PubMed:[18559669](http://www.uniprot.org/citations/18559669)), PubMed:[18829466](http://www.uniprot.org/citations/18829466)), PubMed:[26742426](http://www.uniprot.org/citations/26742426)), PubMed:[28074573](http://www.uniprot.org/citations/28074573)). Positively regulates MAPK signal transduction pathway (PubMed:[28074573](http://www.uniprot.org/citations/28074573)).

Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:[28074573](http://www.uniprot.org/citations/28074573)).

Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity (PubMed:18559669). Dephosphorylates CDC73 (PubMed:26742426). Dephosphorylates SOX9 on tyrosine residues, leading to inactivate SOX9 and promote ossification (By similarity). Dephosphorylates tyrosine-phosphorylated NEDD9/CAS-L (PubMed:19275884).

Cellular Location

Cytoplasm. Nucleus

Tissue Location


Widely expressed, with highest levels in heart, brain, and skeletal muscle.


SH-PTP2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SH-PTP2 Polyclonal Antibody - Images





SH-PTP2 Polyclonal Antibody - Background

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus. Positively regulates MAPK signal transduction pathway (PubMed:28074573). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:28074573). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulation of its RhoA binding activity. Dephosphorylates CDC73 (PubMed:26742426).