

**Skp2 p45 Polyclonal Antibody**  
Catalog # AP72494**Specification****Skp2 p45 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q13309</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

**Skp2 p45 Polyclonal Antibody - Additional Information****Gene ID** 6502**Other Names**

SKP2; FBXL1; S-phase kinase-associated protein 2; Cyclin-A/CDK2-associated protein p45; F-box protein Skp2; F-box/LRR-repeat protein 1; p45skp2

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Skp2 p45 Polyclonal Antibody - Protein Information****Name** SKP2**Synonyms** FBXL1**Function**

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription (PubMed:<a href="http://www.uniprot.org/citations/11931757" target="\_blank">11931757</a>, PubMed:<a href="http://www.uniprot.org/citations/12435635" target="\_blank">12435635</a>, PubMed:<a href="http://www.uniprot.org/citations/12769844" target="\_blank">12769844</a>, PubMed:<a href="http://www.uniprot.org/citations/12840033" target="\_blank">12840033</a>, PubMed:<a href="http://www.uniprot.org/citations/15342634" target="\_blank">15342634</a>, PubMed:<a href="http://www.uniprot.org/citations/15668399" target="\_blank">15668399</a>, PubMed:<a href="http://www.uniprot.org/citations/15949444" target="\_blank">15949444</a>, PubMed:<a href="http://www.uniprot.org/citations/16103164" target="\_blank">16103164</a>, PubMed:<a href="http://www.uniprot.org/citations/16262255" target="\_blank">16262255</a>, PubMed:<a

href="http://www.uniprot.org/citations/16581786" target="\_blank">16581786</a>, PubMed:<a href="http://www.uniprot.org/citations/16951159" target="\_blank">16951159</a>, PubMed:<a href="http://www.uniprot.org/citations/17908926" target="\_blank">17908926</a>, PubMed:<a href="http://www.uniprot.org/citations/17962192" target="\_blank">17962192</a>, PubMed:<a href="http://www.uniprot.org/citations/22464731" target="\_blank">22464731</a>, PubMed:<a href="http://www.uniprot.org/citations/22770219" target="\_blank">22770219</a>, PubMed:<a href="http://www.uniprot.org/citations/32267835" target="\_blank">32267835</a>). Specifically recognizes phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition (By similarity). Degradation of CDKN1B/p27kip also requires CKS1 (By similarity). Recognizes target proteins ORC1, CDT1, RBL2, KMT2A/MLL1, CDK9, RAG2, NBN, FOXO1, UBP43, YTHDF2, and probably MYC, TOB1 and TAL1 (PubMed:<a href="http://www.uniprot.org/citations/11931757" target="\_blank">11931757</a>, PubMed:<a href="http://www.uniprot.org/citations/12435635" target="\_blank">12435635</a>, PubMed:<a href="http://www.uniprot.org/citations/12769844" target="\_blank">12769844</a>, PubMed:<a href="http://www.uniprot.org/citations/12840033" target="\_blank">12840033</a>, PubMed:<a href="http://www.uniprot.org/citations/15342634" target="\_blank">15342634</a>, PubMed:<a href="http://www.uniprot.org/citations/15668399" target="\_blank">15668399</a>, PubMed:<a href="http://www.uniprot.org/citations/15949444" target="\_blank">15949444</a>, PubMed:<a href="http://www.uniprot.org/citations/16103164" target="\_blank">16103164</a>, PubMed:<a href="http://www.uniprot.org/citations/16581786" target="\_blank">16581786</a>, PubMed:<a href="http://www.uniprot.org/citations/16951159" target="\_blank">16951159</a>, PubMed:<a href="http://www.uniprot.org/citations/17908926" target="\_blank">17908926</a>, PubMed:<a href="http://www.uniprot.org/citations/17962192" target="\_blank">17962192</a>, PubMed:<a href="http://www.uniprot.org/citations/22464731" target="\_blank">22464731</a>, PubMed:<a href="http://www.uniprot.org/citations/32267835" target="\_blank">32267835</a>). Degradation of TAL1 also requires STUB1 (PubMed:<a href="http://www.uniprot.org/citations/17962192" target="\_blank">17962192</a>). Recognizes CDKN1A in association with CCNE1 or CCNE2 and CDK2 (PubMed:<a href="http://www.uniprot.org/citations/16262255" target="\_blank">16262255</a>). Promotes ubiquitination and destruction of CDH1 in a CK1-dependent manner, thereby regulating cell migration (PubMed:<a href="http://www.uniprot.org/citations/22770219" target="\_blank">22770219</a>). Following phosphorylation in response to DNA damage, mediates 'Lys-63'-linked ubiquitination of NBN, promoting ATM recruitment to DNA damage sites and DNA repair via homologous recombination (PubMed:<a href="http://www.uniprot.org/citations/22464731" target="\_blank">22464731</a>).

#### Cellular Location


Cytoplasm. Nucleus




#### Skp2 p45 Polyclonal Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Skp2 p45 Polyclonal Antibody - Images









### **Skp2 p45 Polyclonal Antibody - Background**

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription. Specifically recognizes phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition. Degradation of CDKN1B/p27kip also requires CKS1. Recognizes target proteins ORC1, CDT1, RBL2, KMT2A/MLL1, CDK9, RAG2, FOXO1, UBP43, and probably MYC, TOB1 and TAL1. Degradation of TAL1 also requires STUB1. Recognizes CDKN1A in association with CCNE1 or CCNE2 and CDK2. Promotes ubiquitination and destruction of CDH1 in a CK1-Dependent Manner, thereby regulating cell migration.