

TNF-R2 Polyclonal Antibody
Catalog # AP72871**Specification**

TNF-R2 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P20333
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

TNF-R2 Polyclonal Antibody - Additional Information**Gene ID** 7133**Other Names**

TNFRSF1B; TNFBR; TNFR2; Tumor necrosis factor receptor superfamily member 1B; Tumor necrosis factor receptor 2; TNF-R2; Tumor necrosis factor receptor type II; TNF-RII; TNFR-II; p75; p80 TNF-alpha receptor; CD antigen CD120b; Etanercept

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

TNF-R2 Polyclonal Antibody - Protein Information**Name** TNFRSF1B**Synonyms** TNFBR, TNFR2**Function**

Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.

Cellular Location



[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Tumor necrosis factor-binding protein 2]: Secreted

TNF-R2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TNF-R2 Polyclonal Antibody - Images



TNF-R2 Polyclonal Antibody - Background

Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for

homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.