

Osteoglycin Polyclonal Antibody Catalog # AP73477

Specification

Osteoglycin Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P20774
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

Osteoglycin Polyclonal Antibody - Additional Information

Gene ID 4969

Other Names

OGN; OIF; SLRR3A; Mimecan; Osteoglycin; Osteoinductive factor; OIF

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Osteoglycin Polyclonal Antibody - Protein Information

Name OGN

Synonyms OIF, SLRR3A

Function

Induces bone formation in conjunction with TGF-beta-1 or TGF- beta-2.

Cellular Location

Secreted, extracellular space, extracellular matrix {ECO:0000250|UniProtKB:Q8MJF1}

Tissue Location


Bone.

Osteoglycin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Osteoglycin Polyclonal Antibody - Images



Osteoglycin Polyclonal Antibody - Background

Induces bone formation in conjunction with TGF-beta-1 or TGF-beta-2.