

GABAA R α 1 Polyclonal Antibody
Catalog # AP73752**Specification****GABAA R α 1 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	P14867
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

GABAA R α 1 Polyclonal Antibody - Additional Information

Gene ID 2554

Other Names

GABRA1; Gamma-aminobutyric acid receptor subunit alpha-1; GABA(A) receptor subunit alpha-1

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

GABAA R α 1 Polyclonal Antibody - Protein InformationName GABRA1 ([HGNC:4075](#))**Function**

Alpha subunit of the heteropentameric ligand-gated chloride channel gated by Gamma-aminobutyric acid (GABA), a major inhibitory neurotransmitter in the brain (PubMed: [23909897](http://www.uniprot.org/citations/23909897), PubMed: [25489750](http://www.uniprot.org/citations/25489750), PubMed: [29950725](http://www.uniprot.org/citations/29950725), PubMed: [30602789](http://www.uniprot.org/citations/30602789)). GABA-gated chloride channels, also named GABA(A) receptors (GABAAR), consist of five subunits arranged around a central pore and contain GABA active binding site(s) located at the alpha and beta subunit interface(s) (PubMed: [29950725](http://www.uniprot.org/citations/29950725), PubMed: [30602789](http://www.uniprot.org/citations/30602789)). When activated by GABA, GABAARs selectively allow the flow of chloride anions across the cell membrane down their electrochemical gradient (PubMed: [23909897](http://www.uniprot.org/citations/23909897), PubMed: [29950725](http://www.uniprot.org/citations/29950725), PubMed: [30602789](http://www.uniprot.org/citations/30602789)).

Alpha-1/GABRA1-containing GABAARs are largely synaptic (By similarity). Chloride influx into the postsynaptic neuron following GABAAR opening decreases the neuron ability to generate a new action potential, thereby reducing nerve transmission (By similarity). GABAARs containing alpha-1 and beta-2 or -3 subunits exhibit synaptogenic activity; the gamma-2 subunit being necessary but not sufficient to induce rapid synaptic contacts formation (PubMed:23909897, PubMed:25489750). GABAARs function also as histamine receptor where histamine binds at the interface of two neighboring beta subunits and potentiates GABA response (By similarity). GABAARs containing alpha, beta and epsilon subunits also permit spontaneous chloride channel activity while preserving the structural information required for GABA-gated openings (By similarity). Alpha-1-mediated plasticity in the orbitofrontal cortex regulates context-dependent action selection (By similarity). Together with rho subunits, may also control neuronal and glial GABAergic transmission in the cerebellum (By similarity).

Cellular Location


Postsynaptic cell membrane {ECO:0000250|UniProtKB:P08219}; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:P62813}; Multi-pass membrane protein. Note=Mainly located in GABAergic synapses in granule cells, and also in the extrasynaptic membrane at a lower concentration. {ECO:0000250|UniProtKB:P62813}




GABAA R α 1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GABAA R α 1 Polyclonal Antibody - Images





GABAA R α 1 Polyclonal Antibody - Background

Component of the heteropentameric receptor for GABA, the major inhibitory neurotransmitter in the vertebrate brain. Functions also as histamine receptor and mediates cellular responses to histamine. Functions as receptor for diazepam and various anesthetics, such as pentobarbital; these are bound at a separate allosteric effector binding site. Functions as ligand-gated chloride channel (By similarity).