

FBP1 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7385c**Specification**

FBP1 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P09467
Other Accession	Q9OXD6
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36842
Antigen Region	125-156

FBP1 Antibody (Center) - Additional Information**Gene ID** 2203**Other Names**

Fructose-1, 6-bisphosphatase 1, FB Pase 1, D-fructose-1, 6-bisphosphate 1-phosphohydrolase 1, Liver FB Pase, FBP1, FBP

Target/Specificity

This FBP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 125-156 amino acids from the Central region of human FBP1.

DilutionWB~~1:1000
IHC-P~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FBP1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

FBP1 Antibody (Center) - Protein Information**Name** FBP1

Synonyms FBP

Function Catalyzes the hydrolysis of fructose 1,6-bisphosphate to fructose 6-phosphate in the presence of divalent cations, acting as a rate-limiting enzyme in gluconeogenesis. Plays a role in regulating glucose sensing and insulin secretion of pancreatic beta-cells. Appears to modulate glycerol gluconeogenesis in liver. Important regulator of appetite and adiposity; increased expression of the protein in liver after nutrient excess increases circulating satiety hormones and reduces appetite-stimulating neuropeptides and thus seems to provide a feedback mechanism to limit weight gain.

Tissue Location


Expressed in pancreatic islets.

FBP1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


FBP1 Antibody (Center) - Images




Western blot analysis of FBP1 Antibody (Center) (Cat.# AP7385c) in MCF-7 cell line lysates (35ug/lane). FBP1 (arrow) was detected using the purified Pab.



Western blot analysis of FBP1 antibody (Center) (Cat.# AP7385c) in mouse kidney tissue lysates (35ug/lane). FBP1 (arrow) was detected using the purified Pab.



Western blot analysis of FBP1 (arrow) using rabbit polyclonal FBP1 Antibody (Center) (Cat.#AP7385c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the FBP1 gene.



FBP1 Antibody (Center) (Cat.# AP7385c) IHC analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the FBP1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

FBP1 Antibody (Center) - Background

Fructose-1,6-bisphosphatase 1, a gluconeogenesis regulatory enzyme, catalyzes the hydrolysis of

fructose 1,6-bisphosphate to fructose 6-phosphate and inorganic phosphate.
Fructose-1,6-diphosphatase deficiency is associated with hypoglycemia and metabolic acidosis.

FBP1 Antibody (Center) - References

Visinoni,S., Am. J. Physiol. Endocrinol. Metab. 295 (5), E1132-E1141 (2008)
Kebede,M., Diabetes 57 (7), 1887-1895 (2008)