

CTACK Polyclonal Antibody
Catalog # AP73907**Specification**

CTACK Polyclonal Antibody - Product Information

Application	WB
Primary Accession	O9Y4X3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

CTACK Polyclonal Antibody - Additional Information**Gene ID** 10850**Other Names**

CCL27; ILC; SCYA27; C-C motif chemokine 27; CC chemokine ILC; Cutaneous T-cell-attracting chemokine; CTACK; ESkin; IL-11 R-alpha-locus chemokine; Skinkine; Small-inducible cytokine A27

Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CTACK Polyclonal Antibody - Protein Information**Name** CCL27**Synonyms** ILC, SCYA27**Function**

Chemotactic factor that attracts skin-associated memory T- lymphocytes. May play a role in mediating homing of lymphocytes to cutaneous sites. Binds to CCR10.

Cellular Location

Secreted {ECO:0000250|UniProtKB:Q9Z1X0}.

Tissue Location

Testis, thymus, placenta, ovary and skin.

CTACK Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CTACK Polyclonal Antibody - Images





Image not found : 202004/c30669wb42457.jpg





Image not found : 202004/c30669wb42457.jpg

CTACK Polyclonal Antibody - Background

Chemotactic factor that attracts skin-associated memory T-lymphocytes. May play a role in mediating homing of lymphocytes to cutaneous sites. Binds to CCR10.