

**DC-LAMP Polyclonal Antibody**  
Catalog # AP74074**Specification****DC-LAMP Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O9UQV4</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

**DC-LAMP Polyclonal Antibody - Additional Information****Gene ID** 27074**Other Names**

Lysosome-associated membrane glycoprotein 3 (LAMP-3) (Lysosomal-associated membrane protein 3) (DC-lysosome-associated membrane glycoprotein) (DC LAMP) (Protein TSC403) (CD antigen CD208)

**Dilution**

WB~~WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**DC-LAMP Polyclonal Antibody - Protein Information****Name** LAMP3**Synonyms** DCLAMP, TSC403**Function**

Lysosomal membrane glycoprotein which plays a role in the unfolded protein response (UPR) that contributes to protein degradation and cell survival during proteasomal dysfunction (PubMed:<a href="http://www.uniprot.org/citations/25681212" target="\_blank">25681212</a>). Plays a role in the process of fusion of the lysosome with the autophagosome, thereby modulating the autophagic process (PubMed:<a href="http://www.uniprot.org/citations/24434718" target="\_blank">24434718</a>). Promotes hepatocellular lipogenesis through activation of the PI3K/Akt pathway (PubMed:<a href="http://www.uniprot.org/citations/29056532" target="\_blank">29056532</a>). May also play a role in dendritic cell function and in adaptive immunity (PubMed:<a href="http://www.uniprot.org/citations/9768752" target="\_blank">9768752</a>).

**Cellular Location**

Cell surface. Lysosome membrane; Single-pass type I membrane protein. Cytoplasmic vesicle membrane; Single-pass type I membrane protein. Early endosome membrane; Single-pass type I membrane protein. Note=During dendritic cell maturation, detected on cytoplasmic vesicles (the MHC II compartment) that contain MHC II proteins, LAMP1, LAMP2 and LAMP3 (PubMed:9768752). Detected on lysosomes in mature dendritic cells (PubMed:9768752).

#### Tissue Location


Detected in tonsil interdigitating dendritic cells, in spleen, lymph node, Peyer's patches in the small intestine, in thymus medulla and in B-cells (at protein level). Expressed in lymphoid organs and dendritic cells. Expressed in lung. Up-regulated in carcinomas of the esophagus, colon, rectum, ureter, stomach, breast, fallopian tube, thyroid and parotid tissues



#### DC-LAMP Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### DC-LAMP Polyclonal Antibody - Images





### **DC-LAMP Polyclonal Antibody - Background**

May play a role in dendritic cell function and in adaptive immunity.