

**Bob 1 Polyclonal Antibody**  
Catalog # AP74165**Specification**

---

**Bob 1 Polyclonal Antibody - Product Information**

Application	IHC
Primary Accession	<a href="#">Q16633</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

**Bob 1 Polyclonal Antibody - Additional Information****Gene ID** 5450**Other Names**

POU domain class 2-associating factor 1 (B-cell-specific coactivator OBF-1) (BOB-1) (OCA-B) (OCT-binding factor 1)

**Dilution**

IHC~~IHC-p 1:50-200, ELISA 1:10000-20000

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Bob 1 Polyclonal Antibody - Protein Information****Name** POU2AF1 ([HGNC:9211](#))**Function**

Transcriptional coactivator that specifically associates with either POU2F1/OCT1 or POU2F2/OCT2 (PubMed:<a href="http://www.uniprot.org/citations/7859290" target="\_blank">7859290</a>). It boosts the POU2F1/OCT1 mediated promoter activity and to a lesser extent, that of POU2F2/OCT2 (PubMed:<a href="http://www.uniprot.org/citations/7779176" target="\_blank">7779176</a>). It recognizes the POU domains of POU2F1/OCT1 and POU2F2/OCT2 (PubMed:<a href="http://www.uniprot.org/citations/7779176" target="\_blank">7779176</a>). It is essential for the response of B-cells to antigens and required for the formation of germinal centers (PubMed:<a href="http://www.uniprot.org/citations/7623806" target="\_blank">7623806</a>, PubMed:<a href="http://www.uniprot.org/citations/7859290" target="\_blank">7859290</a>). Regulates IL6 expression in B cells as POU2F2/OCT2 coactivator (By similarity).

**Cellular Location**

Nucleus.

**Tissue Location**



B-cell specific (PubMed:7779176, PubMed:7859290). Detected in mainly in spleen, but also in thymus, peripheral blood leukocyte and small intestine (PubMed:7779176, PubMed:7859290)


### **Bob 1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Bob 1 Polyclonal Antibody - Images**





### **Bob 1 Polyclonal Antibody - Background**

Transcriptional coactivator that specifically associates with either OCT1 or OCT2. It boosts the OCT1 mediated promoter activity and to a lesser extent, that of OCT2. It has no intrinsic DNA-binding activity. It recognizes the POU domains of OCT1 and OCT2. It is essential for the response of B-cells to antigens and required for the formation of germinal centers.