

CD70 Polyclonal Antibody
Catalog # AP74214**Specification****CD70 Polyclonal Antibody - Product Information**

Application	IHC
Primary Accession	P32970
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

CD70 Polyclonal Antibody - Additional Information**Gene ID** 970**Other Names**

CD70 antigen (CD27 ligand) (CD27-L) (Tumor necrosis factor ligand superfamily member 7) (CD antigen CD70)

Dilution

IHC~~IHC-p 1:50-200, ELISA 1:10000-20000

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CD70 Polyclonal Antibody - Protein Information**Name** CD70 ([HGNC:11937](#))**Function**

Expressed at the plasma membrane of B cells, it is the ligand of the CD27 receptor which is specifically expressed at the surface of T cells (PubMed: [28011863](http://www.uniprot.org/citations/28011863), PubMed: [28011864](http://www.uniprot.org/citations/28011864), PubMed: [8387892](http://www.uniprot.org/citations/8387892)). The CD70-CD27 signaling pathway mediates antigen-specific T cell activation and expansion which in turn provides immune surveillance of B cells (PubMed: [28011863](http://www.uniprot.org/citations/28011863), PubMed: [28011864](http://www.uniprot.org/citations/28011864)).

Cellular Location



Cell membrane; Single-pass type II membrane protein

CD70 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD70 Polyclonal Antibody - Images



CD70 Polyclonal Antibody - Background

Cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.