

CD275 Polyclonal Antibody
Catalog # AP74221**Specification****CD275 Polyclonal Antibody - Product Information**

Application	IHC
Primary Accession	O75144
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

CD275 Polyclonal Antibody - Additional Information**Gene ID** 23308**Other Names**

ICOS ligand (B7 homolog 2) (B7-H2) (B7-like protein GI50) (B7-related protein 1) (B7RP-1) (CD antigen CD275)

Dilution

IHC~~IHC-p 1:50-200, ELISA 1:10000-20000

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CD275 Polyclonal Antibody - Protein Information**Name** ICOSLG**Function**

Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion (PubMed: [11007762](http://www.uniprot.org/citations/11007762), PubMed: [11023515](http://www.uniprot.org/citations/11023515), PubMed: [30498080](http://www.uniprot.org/citations/30498080)). Also induces B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function (By similarity). In endothelial cells, required for proper neutrophil transmigration in response to chemoattractants, such as CXCL8/IL8 or N-formyl-methionyl peptides (fMLP) (PubMed: [30498080](http://www.uniprot.org/citations/30498080)).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location



Expressed on peripheral blood B-cells and monocytes, as well as on monocyte-derived dendritic cells (at protein level). [Isoform 2]: Detected only in lymph nodes, leukocytes and spleen. Expressed on activated monocytes and dendritic cells.

CD275 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD275 Polyclonal Antibody - Images



CD275 Polyclonal Antibody - Background

Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function

(By similarity).