

MMP-10 Polyclonal Antibody
Catalog # AP74266**Specification**

MMP-10 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P09238
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

MMP-10 Polyclonal Antibody - Additional Information**Gene ID** 4319**Other Names**

Stromelysin-2 (SL-2) (EC 3.4.24.22) (Matrix metalloproteinase-10) (MMP-10) (Transin-2)

Dilution

WB~~WB 1:500-2000, ELISA 1:10000-20000

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

MMP-10 Polyclonal Antibody - Protein Information**Name** MMP10**Synonyms** STMY2**Function**

Can degrade fibronectin, gelatins of type I, III, IV, and V; weakly collagens III, IV, and V. Activates procollagenase.

Cellular Location

Secreted, extracellular space, extracellular matrix



MMP-10 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MMP-10 Polyclonal Antibody - Images



MMP-10 Polyclonal Antibody - Background

Can degrade fibronectin, gelatins of type I, III, IV, and V; weakly collagens III, IV, and V. Activates procollagenase.