

VTN Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7462b**Specification**

VTN Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P04004
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	54306
Antigen Region	352-379

VTN Antibody (C-term) - Additional Information**Gene ID** 7448**Other Names**

Vitronectin, VN, S-protein, Serum-spreading factor, V75, Vitronectin V65 subunit, Vitronectin V10 subunit, Somatomedin-B, VTN

Target/Specificity

This VTN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 352-379 amino acids from the C-terminal region of human VTN.

DilutionWB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

VTN Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

VTN Antibody (C-term) - Protein Information**Name** VTN

Function Vitronectin is a cell adhesion and spreading factor found in serum and tissues. Vitronectin interact with glycosaminoglycans and proteoglycans. Is recognized by certain members of the integrin family and serves as a cell-to-substrate adhesion molecule. Inhibitor of the membrane-damaging effect of the terminal cytolytic complement pathway.

Cellular Location

Secreted, extracellular space

Tissue Location


Expressed in the retina pigment epithelium (at protein level) (PubMed:25136834). Expressed in plasma (at protein level) (PubMed:2448300). Expressed in serum (at protein level) (PubMed:29567995).

VTN Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


VTN Antibody (C-term) - Images



Western blot analysis of VTN Antibody (C-term) (Cat.#AP7462b) in NCI-H460(lane 1), HepG2(lane 2) cell line lysates (35ug/lane). VTN (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma with VTN Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of NCI-H292 cells using VTN Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

VTN Antibody (C-term) - Background

VTN is a member of the pexin family. This protein is found in serum and tissues and promotes cell adhesion and spreading, inhibits the membrane-damaging effect of the terminal cytolytic complement pathway, and binds to several serpin serine protease inhibitors. The protein is a secreted protein and exists in either a single chain form or a clipped, two chain form held together by a disulfide bond.

VTN Antibody (C-term) - References

- Jenne D.E., Stanley K.K.EMBO J. 4:3153-3157(1985)
- Sigurdardottir O., Wiman B.Biochim. Acta 1208:104-110(1994)
- Seiffert D., Loskutoff D.J.J. Biol. Chem. 266:2824-2830(1991)