

**RabbitAnti-CD9 antibody**

CD9

Catalog # AP74804

**Specification****RabbitAnti-CD9 antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P21926</a>
Reactivity	Human, Mouse
Predicted	Human, Mouse
Host	Rabbit
Clonality	Monoclonal
Calculated MW	25kDa KDa

**RabbitAnti-CD9 antibody - Additional Information**

Gene ID 928

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**RabbitAnti-CD9 antibody - Protein Information**

Name CD9 {ECO:0000303|PubMed:1840589, ECO:0000312|HGNC:HGNC:1709}

**Function**

Integral membrane protein associated with integrins, which regulates different processes, such as sperm-egg fusion, platelet activation and aggregation, and cell adhesion (PubMed:[14575715](http://www.uniprot.org/citations/14575715), PubMed:[18541721](http://www.uniprot.org/citations/18541721), PubMed:[8478605](http://www.uniprot.org/citations/8478605)). Present at the cell surface of oocytes and plays a key role in sperm-egg fusion, possibly by organizing multiprotein complexes and the morphology of the membrane required for the fusion (By similarity). In myoblasts, associates with CD81 and PTGFRN and inhibits myotube fusion during muscle regeneration (By similarity). In macrophages, associates with CD81 and beta-1 and beta-2 integrins, and prevents macrophage fusion into multinucleated giant cells specialized in ingesting complement-opsonized large particles (PubMed:[12796480](http://www.uniprot.org/citations/12796480)). Also prevents the fusion between mononuclear cell progenitors into osteoclasts in charge of bone resorption (By similarity). Acts as a receptor for PSG17 (By similarity). Involved in platelet activation and aggregation (PubMed:[18541721](http://www.uniprot.org/citations/18541721)). Regulates paranodal junction formation (By similarity). Involved in cell adhesion, cell motility and tumor metastasis (PubMed:[7511626](http://www.uniprot.org/citations/7511626), PubMed:[7511626](http://www.uniprot.org/citations/7511626)).

href="http://www.uniprot.org/citations/8478605" target="\_blank">8478605</a>).

### Cellular Location

Cell membrane; Multi-pass membrane protein. Membrane; Multi-pass membrane protein. Secreted, extracellular exosome {ECO:0000250|UniProtKB:P40240}. Note=Present at the cell surface of oocytes. Accumulates in the adhesion area between the sperm and egg following interaction between IZUMO1 and its receptor IZUMO1R/JUNO {ECO:0000250|UniProtKB:P40240}

### Tissue Location


Detected in platelets (at protein level) (PubMed:19640571). Expressed by a variety of hematopoietic and epithelial cells (PubMed:19640571).

### RabbitAnti-CD9 antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### RabbitAnti-CD9 antibody - Images




Sample: Lane 1: Kidney (Mouse) Lysate at 40 ug Lane 2: Testis (Mouse) Lysate at 40 ug Lane 3: Heart (Mouse) Lysate at 40 ug Lane 4: Cerebrum (Mouse) Lysate at 40 ug Lane 5: Huvec (Human) Cell Lysate at 30 ug Lane 6: A431 (Human) Cell Lysate at 30 ug Lane 7: SW480 (Human) Cell Lysate at 30 ug Primary: Anti-CD9 (bsm-52264R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 24 kD Observed band size: 24 kD




Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CD9) Monoclonal Antibody, Unconjugated (bsm-52264R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CD9) Monoclonal Antibody, Unconjugated (bsm-52264R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human laryngeal carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CD9) Monoclonal Antibody, Unconjugated (bsm-52264R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CD9) Monoclonal Antibody, Unconjugated (bsm-52264R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.