

MCM6 Rabbit mAb
Catalog # AP74817**Specification****MCM6 Rabbit mAb - Product Information**

Application	WB, IHC, IF
Primary Accession	Q14566
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	92889

MCM6 Rabbit mAb - Additional Information

Gene ID 4175

Other Names
MCM6**Dilution**
WB~~1/500-1/1000
IHC~~1/50-1/100
IF~~1/50-1/200**Format**
Liquid**MCM6 Rabbit mAb - Protein Information**Name MCM6 ([HGNC:6949](#))**Function**

Acts as a component of the MCM2-7 complex (MCM complex) which is the replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Core component of CDC45-MCM-GINS (CMG) helicase, the molecular machine that unwinds template DNA during replication, and around which the replisome is built (PubMed:16899510, PubMed:32453425, PubMed:34694004, PubMed:34700328, PubMed:35585232, PubMed:9305914). The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity (PubMed:32453425).

Cellular Location

Nucleus. Chromosome. Note= Binds to chromatin during G1 and detaches from it during S phase.

MCM6 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MCM6 Rabbit mAb - Images



