

**Toll-Like Receptor 2 Rabbit mAb**  
Catalog # AP75005**Specification****Toll-Like Receptor 2 Rabbit mAb - Product Information**

Application	WB, IHC, IF
Primary Accession	<a href="#">O9OUN7</a>
Reactivity	Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	89449

**Toll-Like Receptor 2 Rabbit mAb - Additional Information**

Gene ID 24088

**Other Names**

Tlr2

**Dilution**

WB~~1/500-1/1000

IHC~~1/50-1/100

IF~~1/50-1/200

**Format**

Liquid

**Toll-Like Receptor 2 Rabbit mAb - Protein Information**

Name Tlr2

**Function**

Cooperates with LY96 to mediate the innate immune response to bacterial lipoproteins and other microbial cell wall components. Cooperates with TLR1 or TLR6 to mediate the innate immune response to bacterial lipoproteins or lipopeptides. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (By similarity) (PubMed:<a href="http://www.uniprot.org/citations/15690042" target="\_blank">15690042</a>). May also promote apoptosis in response to lipoproteins (By similarity). Forms activation clusters composed of several receptors depending on the ligand, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway. Forms the cluster TLR2:TLR6:CD14:CD36 in response to diacylated lipopeptides and TLR2:TLR1:CD14 in response to triacylated lipopeptides (By similarity). Recognizes M.tuberculosis major T-antigen EsxA (ESAT-6) which inhibits downstream MYD88-dependent signaling (PubMed:<a href="http://www.uniprot.org/citations/17486091" target="\_blank">17486091</a>). Acts as the major receptor for M.tuberculosis lipoproteins LprA, LprG, LpqH and PhoS1 (pstS1), in conjunction with TLR1 and for some but not all lipoproteins CD14 and/or CD36. The lipoproteins act as agonists to modulate antigen presenting cell functions in response to the pathogen (PubMed:<a href="http://www.uniprot.org/citations/19362712" target="\_blank">19362712</a>). Recombinant

MPT83 from *M.tuberculosis* stimulates secretion of cytokines (TNF-alpha, IL-6 and IL-12p40) by mouse macrophage cell lines in a TLR2-dependent fashion, which leads to increased host innate immunity responses against the bacterium (PubMed:<a href="http://www.uniprot.org/citations/22174456" target="\_blank">22174456</a>). Lung macrophages which express low levels of TLR2 respond poorly to stimulation by *M.tuberculosis* LpqH (PubMed:<a href="http://www.uniprot.org/citations/19362712" target="\_blank">19362712</a>). Required for normal uptake of *M.tuberculosis*, a process that is inhibited by *M.tuberculosis* LppM (PubMed:<a href="http://www.uniprot.org/citations/27220037" target="\_blank">27220037</a>). Interacts with TICAM2 (By similarity).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Cytoplasmic vesicle, phagosome membrane; Single-pass type I membrane protein. Membrane raft {ECO:0000250|UniProtKB:O60603}. Note=Does not reside in lipid rafts before stimulation but accumulates increasingly in the raft upon the presence of the microbial ligand. In response to diacylated lipoproteins, TLR2:TLR6 heterodimers are recruited in lipid rafts, this recruitment determine the intracellular targeting to the Golgi apparatus. Triacylated lipoproteins induce the same mechanism for TLR2:TLR1 heterodimers. {ECO:0000250|UniProtKB:O60603}

#### Tissue Location

Detected in a macrophage cell line, smooth muscle, lung, spleen, thymus, brain and adipose tissue. Cell surface expression detected in lung alveolar macrophages, dendritic macrophages and at lower levels in lung macrophages (at protein level) (PubMed:19362712)

#### Toll-Like Receptor 2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Toll-Like Receptor 2 Rabbit mAb - Images



