

**alpha Synuclein Rabbit mAb**  
Catalog # AP75067**Specification****alpha Synuclein Rabbit mAb - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P37840</a>
Reactivity	Human, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	14460

**alpha Synuclein Rabbit mAb - Additional Information**

Gene ID 6622

**Other Names**

SNCA

**Dilution**

WB~~1/500-1/1000

IHC~~1/50-1/100

**Format**

Liquid

**alpha Synuclein Rabbit mAb - Protein Information**

Name SNCA

Synonyms NACP, PARK1

**Function**

Neuronal protein that plays several roles in synaptic activity such as regulation of synaptic vesicle trafficking and subsequent neurotransmitter release (PubMed: [20798282](http://www.uniprot.org/citations/20798282), PubMed: [26442590](http://www.uniprot.org/citations/26442590), PubMed: [28288128](http://www.uniprot.org/citations/28288128), PubMed: [30404828](http://www.uniprot.org/citations/30404828)). Participates as a monomer in synaptic vesicle exocytosis by enhancing vesicle priming, fusion and dilation of exocytotic fusion pores (PubMed: [28288128](http://www.uniprot.org/citations/28288128), PubMed: [30404828](http://www.uniprot.org/citations/30404828)). Mechanistically, acts by increasing local Ca(2+) release from microdomains which is essential for the enhancement of ATP-induced exocytosis (PubMed: [30404828](http://www.uniprot.org/citations/30404828)). Acts also as a molecular chaperone in its multimeric membrane-bound state, assisting in the folding of synaptic fusion components called SNAREs (Soluble NSF Attachment Protein REceptors) at presynaptic plasma membrane in conjunction with cysteine string protein-alpha/DNAJC5

(PubMed:<a href="http://www.uniprot.org/citations/20798282" target="\_blank">20798282</a>). This chaperone activity is important to sustain normal SNARE-complex assembly during aging (PubMed:<a href="http://www.uniprot.org/citations/20798282" target="\_blank">20798282</a>). Also plays a role in the regulation of the dopamine neurotransmission by associating with the dopamine transporter (DAT1) and thereby modulating its activity (PubMed:<a href="http://www.uniprot.org/citations/26442590" target="\_blank">26442590</a>).

#### Cellular Location

Cytoplasm. Membrane. Nucleus. Synapse Secreted. Cell projection, axon {ECO:0000250|UniProtKB:O55042}. Note=Membrane-bound in dopaminergic neurons (PubMed:15282274). Expressed and colocalized with SEPTIN4 in dopaminergic axon terminals, especially at the varicosities (By similarity). {ECO:0000250|UniProtKB:O55042, ECO:0000269|PubMed:15282274}

#### Tissue Location

Highly expressed in presynaptic terminals in the central nervous system. Expressed principally in brain

### alpha Synuclein Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### alpha Synuclein Rabbit mAb - Images



