

CD3G Rabbit mAb
Catalog # AP75230**Specification**

CD3G Rabbit mAb - Product Information

Application	WB, IF
Primary Accession	P09693
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	20469

CD3G Rabbit mAb - Additional Information**Gene ID** 917**Other Names**
CD3G**Dilution**
WB~~1/500-1/1000
IF~~1/50-1/200**Format**
Liquid**CD3G Rabbit mAb - Protein Information****Name** CD3G**Synonyms** T3G**Function**

Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed: [2470098](http://www.uniprot.org/citations/2470098)). In addition to this role of signal transduction in T-cell activation, CD3G plays an essential role in the dynamic regulation of TCR expression at the cell surface (PubMed: [8187769](http://www.uniprot.org/citations/8187769)). Indeed, constitutive TCR cycling is dependent on the di-leucine-based (diL) receptor-sorting motif present in CD3G.

Cellular Location

Cell membrane; Single-pass type I membrane protein

CD3G Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD3G Rabbit mAb - Images

