

**EpCAM Rabbit mAb**  
**Catalog # AP75405****Specification**

---

**EpCAM Rabbit mAb - Product Information**

Application	WB, IHC, IF, IP
Primary Accession	<a href="#">P16422</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	34932

**EpCAM Rabbit mAb - Additional Information****Gene ID** 4072**Other Names**  
EPCAM**Dilution**  
WB~~1/500-1/1000  
IHC~~1/50-1/100  
IF~~1/50-1/200  
IP~~1/20**Format**  
Liquid**EpCAM Rabbit mAb - Protein Information****Name** EPCAM**Synonyms** GA733-2, M1S2, M4S1, MIC18, TACSTD1, TRO**Function**

May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.

**Cellular Location**

Lateral cell membrane; Single-pass type I membrane protein. Cell junction, tight junction.  
Note=Colocalizes with CLDN7 at the lateral cell membrane and tight junction

**Tissue Location**

Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC) Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on

the surface of adenocarcinoma.

### EpCAM Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### EpCAM Rabbit mAb - Images



