

**Fibrillarlin Rabbit mAb**  
Catalog # AP75438**Specification****Fibrillarlin Rabbit mAb - Product Information**

Application	WB
Primary Accession	<a href="#">P22087</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	33784

**Fibrillarlin Rabbit mAb - Additional Information**

Gene ID 2091

**Other Names**

FBL

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**Fibrillarlin Rabbit mAb - Protein Information**Name FBL ([HGNC:3599](#))

Synonyms FIB1, FLRN

**Function**

S-adenosyl-L-methionine-dependent methyltransferase that has the ability to methylate both RNAs and proteins (PubMed: [24352239](http://www.uniprot.org/citations/24352239), PubMed: [30540930](http://www.uniprot.org/citations/30540930), PubMed: [32017898](http://www.uniprot.org/citations/32017898)). Involved in pre-rRNA processing by catalyzing the site-specific 2'-hydroxyl methylation of ribose moieties in pre-ribosomal RNA (PubMed: [30540930](http://www.uniprot.org/citations/30540930)). Site specificity is provided by a guide RNA that base pairs with the substrate (By similarity). Methylation occurs at a characteristic distance from the sequence involved in base pairing with the guide RNA (By similarity). Probably catalyzes 2'-O-methylation of U6 snRNAs in box C/D RNP complexes (PubMed: [32017898](http://www.uniprot.org/citations/32017898)). U6 snRNA 2'-O-methylation is required for mRNA splicing fidelity (PubMed: [32017898](http://www.uniprot.org/citations/32017898)). Also acts as a protein methyltransferase by mediating methylation of 'Gln-105' of histone H2A (H2AQ104me), a modification that impairs binding of the FACT complex and is specifically present at 35S ribosomal DNA locus (PubMed: [32017898](#))

[24352239](http://www.uniprot.org/citations/24352239), PubMed:<[30540930](http://www.uniprot.org/citations/30540930)>). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed:<[34516797](http://www.uniprot.org/citations/34516797)>).

#### Cellular Location

Nucleus, nucleolus. Nucleus, nucleoplasm {ECO:0000250|UniProtKB:P35550}. Note=Fibrillar region of the nucleolus

#### Fibrillarin Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Fibrillarin Rabbit mAb - Images

