

GRP78 BiP Rabbit mAb
Catalog # AP75519**Specification****GRP78 BiP Rabbit mAb - Product Information**

Application	WB, IHC
Primary Accession	P11021
Reactivity	Human, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	72333

GRP78 BiP Rabbit mAb - Additional Information

Gene ID 3309

Other Names
HSPA5**Dilution**
WB~~1/500-1/1000
IHC~~1/50-1/100**Format**
Liquid**GRP78 BiP Rabbit mAb - Protein Information****Name** HSPA5 ([HGNC:5238](#))**Function**

Endoplasmic reticulum chaperone that plays a key role in protein folding and quality control in the endoplasmic reticulum lumen (PubMed: [2294010](http://www.uniprot.org/citations/2294010), PubMed: [23769672](http://www.uniprot.org/citations/23769672), PubMed: [23990668](http://www.uniprot.org/citations/23990668), PubMed: [28332555](http://www.uniprot.org/citations/28332555)). Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10/ERdj5, probably to facilitate the release of DNAJC10/ERdj5 from its substrate (By similarity). Acts as a key repressor of the ERN1/IRE1-mediated unfolded protein response (UPR) (PubMed: [1550958](http://www.uniprot.org/citations/1550958), PubMed: [19538957](http://www.uniprot.org/citations/19538957)). In the unstressed endoplasmic reticulum, recruited by DNAJB9/ERdj4 to the luminal region of ERN1/IRE1, leading to disrupt the dimerization of ERN1/IRE1, thereby inactivating ERN1/IRE1 (By similarity). Accumulation of misfolded protein in the endoplasmic reticulum causes release of HSPA5/BiP from ERN1/IRE1, allowing homodimerization and subsequent activation of ERN1/IRE1 (By similarity). Plays an auxiliary role in post-translational transport of small presecretory proteins across endoplasmic reticulum (ER). May function as an allosteric modulator for SEC61 channel-forming

translocon complex, likely cooperating with SEC62 to enable the productive insertion of these precursors into SEC61 channel. Appears to specifically regulate translocation of precursors having inhibitory residues in their mature region that weaken channel gating. May also play a role in apoptosis and cell proliferation (PubMed:26045166).

Cellular Location

Endoplasmic reticulum lumen. Melanosome. Cytoplasm {ECO:0000250|UniProtKB:P20029}. Cell surface Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545). Localizes to the cell surface of epithelial cells in response to high levels of free iron (PubMed:20484814, PubMed:24355926, PubMed:27159390)

GRP78 BiP Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRP78 BiP Rabbit mAb - Images



