

**HMG4 Rabbit mAb**  
Catalog # AP75553**Specification**

---

**HMG4 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O15347</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>22980</b>

**HMG4 Rabbit mAb - Additional Information****Gene ID** 3149**Other Names**  
HMGB3**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**HMG4 Rabbit mAb - Protein Information****Name** HMGB3**Synonyms** HMG2A, HMG4**Function**

Multifunctional protein with various roles in different cellular compartments. May act in a redox sensitive manner. Associates with chromatin and binds DNA with a preference for non-canonical DNA structures such as single-stranded DNA. Can bend DNA and enhance DNA flexibility by looping thus providing a mechanism to promote activities on various gene promoters (By similarity). Proposed to be involved in the innate immune response to nucleic acids by acting as a cytoplasmic promiscuous immunogenic DNA/RNA sensor (By similarity). Negatively regulates B-cell and myeloid cell differentiation. In hematopoietic stem cells may regulate the balance between self-renewal and differentiation. Involved in negative regulation of canonical Wnt signaling (By similarity).

**Cellular Location**

Nucleus {ECO:0000250|UniProtKB:P40618, ECO:0000255|PROSITE-ProRule:PRU00267}.  
Chromosome Cytoplasm {ECO:0000250|UniProtKB:O54879}

**Tissue Location**

Expressed predominantly in placenta.

## HMG4 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## HMG4 Rabbit mAb - Images

