

## HP1 gamma Rabbit mAb

**Catalog # AP75570** 

# **Specification**

# **HP1** gamma Rabbit mAb - Product Information

Application WB, IHC Primary Accession P23198

Reactivity Human, Mouse, Rat

Host Rabb

Clonality Monoclonal Antibody

Calculated MW 20855

## **HP1** gamma Rabbit mAb - Additional Information

**Other Names** 

Cbx3

**Dilution** 

WB~~1/500-1/1000 IHC~~1/50-1/100

**Format** 

Liquid

# HP1 gamma Rabbit mAb - Protein Information

### Name Cbx3

#### **Function**

Component of heterochromatin. Recognizes and binds histone H3 tails methylated at 'Lys-9', leading to epigenetic repression. Probably involved in the repression of many genes located in euchromatin, such as E2F1, MYC and CDC25A. Involved in the formation of functional kinetochore through interaction with MIS12 complex proteins. Contributes to the conversion of local chromatin to a heterochromatin- like repressive state through H3 'Lys-9' trimethylation, mediates the recruitment of the methyltransferases SUV39H1 and/or SUV39H2 by the PER complex to the E-box elements of the circadian target genes such as PER2 itself or PER1 (PubMed:<a href="http://www.uniprot.org/citations/24413057" target="\_blank">24413057</a>). Mediates the recruitment of NIPBL to sites of DNA damage at double-strand breaks (DSBs) (By similarity).

### **Cellular Location**

Nucleus. Note=May be associated with microtubules and mitotic poles during mitosis (Potential). Associates with euchromatin and is largely excluded from constitutive heterochromatin.

### **HP1** gamma Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.



- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# HP1 gamma Rabbit mAb - Images



