

## **ACAT1 Antibody (C-term)**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7560b

### **Specification**

## ACAT1 Antibody (C-term) - Product Information

Application WB, IHC-P,E
Primary Accession P24752
Other Accession O8HXY6

Reactivity Human, Mouse

Predicted Monkey
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 45200
Antigen Region 296-329

# **ACAT1 Antibody (C-term) - Additional Information**

#### Gene ID 38

### **Other Names**

Acetyl-CoA acetyltransferase, mitochondrial, Acetoacetyl-CoA thiolase, T2, ACAT1, ACAT, MAT

### Target/Specificity

This ACAT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 296-329 amino acids from the C-terminal region of human ACAT1.

### **Dilution**

WB~~1:1000 IHC-P~~1:10~50

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

#### Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

ACAT1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# **ACAT1 Antibody (C-term) - Protein Information**

### Name ACAT1





Synonyms ACAT, MAT

**Function** This is one of the enzymes that catalyzes the last step of the mitochondrial beta-oxidation pathway, an aerobic process breaking down fatty acids into acetyl-CoA (PubMed:1715688, PubMed:7728148, PubMed:9744475). Using free coenzyme A/CoA, catalyzes the thiolytic cleavage of medium- to long-chain 3-oxoacyl-CoAs into acetyl-CoA and a fatty acyl-CoA shortened by two carbon atoms (PubMed:1715688, PubMed:7728148, PubMed:9744475). The activity of the enzyme is reversible and it can also catalyze the condensation of two acetyl-CoA molecules into acetoacetyl-CoA (PubMed:17371050). Thereby, it plays a major role in ketone body metabolism (PubMed:1715688, PubMed:17371050, PubMed:7728148, PubMed:9744475).


**Cellular Location**Mitochondrion.

## **ACAT1 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## ACAT1 Antibody (C-term) - Images




Western blot analysis of anti-ACAT1 Pab (Cat.#AP7560b) in SK-BR-3 cell line lysates (35ug/lane).ACAT1 (arrow) was detected using the purified Pab.






All lanes : Anti-ACAT1 Antibody (C-term) at 1:2000 dilution Lane 1: SW620 whole cell lysates Lane 2: mouse liver lysates Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 45. 2 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes: Anti-ACAT1 Antibody (C-term) at 1:1000 dilution Lane 1: SK-BR-3 whole cell lysates Lane 2: HepG2 whole cell lysates Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size: 45. 2 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





Tel: 858.875.1900 Fax: 858.875.1999

Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with ACAT1 antibody (C-term) (Cat.#AP7560b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

## ACAT1 Antibody (C-term) - Background

ACAT1 is a mitochondrially localized enzyme that catalyzes the reversible formation of acetoacetyl-CoA from two molecules of acetyl-CoA. Defects in the gene encoding ACAT1 are associated with the alpha-methylacetoaceticaciduria disorder, an inborn error of isoleucine catabolism characterized by urinary excretion of 2-methyl-3-hydroxybutyric acid, 2-methylacetoacetic acid, tiglylglycine, and butanone.

### ACAT1 Antibody (C-term) - References

Locke, J.A., Prostate 68 (1), 20-33 (2008) Guo, Z.Y., Biochemistry 46 (35), 10063-10071 (2007) Haapalainen, A.M., Biochemistry 46 (14), 4305-4321 (2007)