

**Mre11 Rabbit mAb**  
Catalog # AP75736**Specification****Mre11 Rabbit mAb - Product Information**

Application	<b>WB, IHC</b>
Primary Accession	<a href="#">P49959</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>80593</b>

**Mre11 Rabbit mAb - Additional Information**

Gene ID 4361

**Other Names**  
MRE11**Dilution**  
WB~~1/500-1/1000  
IHC~~1/50-1/100**Format**  
Liquid**Mre11 Rabbit mAb - Protein Information****Name** MRE11 {ECO:0000303|PubMed:8530104, ECO:0000312|HGNC:HGNC:7230}**Function**

Core component of the MRN complex, which plays a central role in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity and meiosis (PubMed:<a href="http://www.uniprot.org/citations/11741547" target="\_blank">11741547</a>, PubMed:<a href="http://www.uniprot.org/citations/14657032" target="\_blank">14657032</a>, PubMed:<a href="http://www.uniprot.org/citations/22078559" target="\_blank">22078559</a>, PubMed:<a href="http://www.uniprot.org/citations/23080121" target="\_blank">23080121</a>, PubMed:<a href="http://www.uniprot.org/citations/24316220" target="\_blank">24316220</a>, PubMed:<a href="http://www.uniprot.org/citations/26240375" target="\_blank">26240375</a>, PubMed:<a href="http://www.uniprot.org/citations/27889449" target="\_blank">27889449</a>, PubMed:<a href="http://www.uniprot.org/citations/28867292" target="\_blank">28867292</a>, PubMed:<a href="http://www.uniprot.org/citations/29670289" target="\_blank">29670289</a>, PubMed:<a href="http://www.uniprot.org/citations/30464262" target="\_blank">30464262</a>, PubMed:<a href="http://www.uniprot.org/citations/30612738" target="\_blank">30612738</a>, PubMed:<a href="http://www.uniprot.org/citations/31353207" target="\_blank">31353207</a>, PubMed:<a href="http://www.uniprot.org/citations/37696958" target="\_blank">37696958</a>, PubMed:<a href="http://www.uniprot.org/citations/38128537" target="\_blank">38128537</a>, PubMed:<a href="http://www.uniprot.org/citations/9590181" target="\_blank">9590181</a>, PubMed:<a

href="http://www.uniprot.org/citations/9651580" target="\_blank">9651580</a>, PubMed:<a href="http://www.uniprot.org/citations/9705271" target="\_blank">9705271</a>). The MRN complex is involved in the repair of DNA double-strand breaks (DSBs) via homologous recombination (HR), an error-free mechanism which primarily occurs during S and G2 phases (PubMed:<a href="http://www.uniprot.org/citations/24316220" target="\_blank">24316220</a>, PubMed:<a href="http://www.uniprot.org/citations/28867292" target="\_blank">28867292</a>, PubMed:<a href="http://www.uniprot.org/citations/31353207" target="\_blank">31353207</a>, PubMed:<a href="http://www.uniprot.org/citations/38128537" target="\_blank">38128537</a>). The complex (1) mediates the end resection of damaged DNA, which generates proper single-stranded DNA, a key initial steps in HR, and is (2) required for the recruitment of other repair factors and efficient activation of ATM and ATR upon DNA damage (PubMed:<a href="http://www.uniprot.org/citations/24316220" target="\_blank">24316220</a>, PubMed:<a href="http://www.uniprot.org/citations/27889449" target="\_blank">27889449</a>, PubMed:<a href="http://www.uniprot.org/citations/28867292" target="\_blank">28867292</a>, PubMed:<a href="http://www.uniprot.org/citations/36050397" target="\_blank">36050397</a>, PubMed:<a href="http://www.uniprot.org/citations/38128537" target="\_blank">38128537</a>). Within the MRN complex, MRE11 possesses both single-strand endonuclease activity and double-strand-specific 3'-5' exonuclease activity (PubMed:<a href="http://www.uniprot.org/citations/11741547" target="\_blank">11741547</a>, PubMed:<a href="http://www.uniprot.org/citations/22078559" target="\_blank">22078559</a>, PubMed:<a href="http://www.uniprot.org/citations/24316220" target="\_blank">24316220</a>, PubMed:<a href="http://www.uniprot.org/citations/26240375" target="\_blank">26240375</a>, PubMed:<a href="http://www.uniprot.org/citations/27889449" target="\_blank">27889449</a>, PubMed:<a href="http://www.uniprot.org/citations/29670289" target="\_blank">29670289</a>, PubMed:<a href="http://www.uniprot.org/citations/31353207" target="\_blank">31353207</a>, PubMed:<a href="http://www.uniprot.org/citations/36563124" target="\_blank">36563124</a>, PubMed:<a href="http://www.uniprot.org/citations/9590181" target="\_blank">9590181</a>, PubMed:<a href="http://www.uniprot.org/citations/9651580" target="\_blank">9651580</a>, PubMed:<a href="http://www.uniprot.org/citations/9705271" target="\_blank">9705271</a>). After DSBs, MRE11 is loaded onto DSBs sites and cleaves DNA by cooperating with RBBP8/CtIP to initiate end resection (PubMed:<a href="http://www.uniprot.org/citations/27814491" target="\_blank">27814491</a>, PubMed:<a href="http://www.uniprot.org/citations/27889449" target="\_blank">27889449</a>, PubMed:<a href="http://www.uniprot.org/citations/30787182" target="\_blank">30787182</a>). MRE11 first endonucleolytically cleaves the 5' strand at DNA DSB ends to prevent non-homologous end joining (NHEJ) and license HR (PubMed:<a href="http://www.uniprot.org/citations/24316220" target="\_blank">24316220</a>). It then generates a single-stranded DNA gap via 3' to 5' exonucleolytic degradation to create entry sites for EXO1- and DNA2-mediated 5' to 3' long-range resection, which is required for single-strand invasion and recombination (PubMed:<a href="http://www.uniprot.org/citations/24316220" target="\_blank">24316220</a>, PubMed:<a href="http://www.uniprot.org/citations/28867292" target="\_blank">28867292</a>). RBBP8/CtIP specifically promotes the endonuclease activity of MRE11 to clear protein-DNA adducts and generate clean double-strand break ends (PubMed:<a href="http://www.uniprot.org/citations/27814491" target="\_blank">27814491</a>, PubMed:<a href="http://www.uniprot.org/citations/27889449" target="\_blank">27889449</a>, PubMed:<a href="http://www.uniprot.org/citations/30787182" target="\_blank">30787182</a>). The MRN complex is also required for DNA damage signaling via activation of the ATM and ATR kinases: the nuclease activity of MRE11 is not required to activate ATM and ATR (PubMed:<a href="http://www.uniprot.org/citations/14657032" target="\_blank">14657032</a>, PubMed:<a href="http://www.uniprot.org/citations/15064416" target="\_blank">15064416</a>, PubMed:<a href="http://www.uniprot.org/citations/15790808" target="\_blank">15790808</a>, PubMed:<a href="http://www.uniprot.org/citations/16622404" target="\_blank">16622404</a>). The MRN complex is also required for the processing of R-loops (PubMed:<a href="http://www.uniprot.org/citations/31537797" target="\_blank">31537797</a>). The MRN complex is involved in the activation of the cGAS-STING pathway induced by DNA damage during tumorigenesis: the MRN complex acts by displacing CGAS from nucleosome sequestration, thereby activating it (By similarity). In telomeres the MRN complex may modulate t-loop formation (PubMed:<a href="http://www.uniprot.org/citations/10888888" target="\_blank">10888888</a>).

### Cellular Location

Nucleus. Chromosome. Chromosome, telomere Note=Localizes to DNA double-strand breaks (DSBs)

### Mre11 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Mre11 Rabbit mAb - Images



