

**NXF1 Rabbit mAb**  
Catalog # AP75837**Specification****NXF1 Rabbit mAb - Product Information**

Application	WB, IHC, IF
Primary Accession	<a href="#">Q9UBU9</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	70182

**NXF1 Rabbit mAb - Additional Information**

Gene ID 10482

**Other Names**

NXF1

**Dilution**

WB~~1/500-1/1000

IHC~~1/50-1/100

IF~~1/50-1/200

**Format**

Liquid

**NXF1 Rabbit mAb - Protein Information**

Name NXF1

Synonyms TAP

**Function**

Involved in the nuclear export of mRNA species bearing retroviral constitutive transport elements (CTE) and in the export of mRNA from the nucleus to the cytoplasm (TAP/NFX1 pathway) (PubMed: [10924507](http://www.uniprot.org/citations/10924507)). The NXF1-NXT1 heterodimer is involved in the export of HSP70 mRNA in conjunction with ALYREF/THOC4 and THOC5 components of the TREX complex (PubMed: [18364396](http://www.uniprot.org/citations/18364396), PubMed: [19165146](http://www.uniprot.org/citations/19165146), PubMed: [9660949](http://www.uniprot.org/citations/9660949)). ALYREF/THOC4-bound mRNA is thought to be transferred to the NXF1-NXT1 heterodimer for export (PubMed: [18364396](http://www.uniprot.org/citations/18364396), PubMed: [19165146](http://www.uniprot.org/citations/19165146), PubMed: [9660949](http://www.uniprot.org/citations/9660949)). Also involved in nuclear export of m6A-containing mRNAs: interaction between SRSF3 and YTHDC1 facilitates m6A-containing mRNA-binding to both SRSF3 and NXF1, promoting mRNA nuclear

export (PubMed:<a href="http://www.uniprot.org/citations/28984244" target="\_blank">28984244</a>).

#### Cellular Location

Nucleus. Nucleus, nucleoplasm Nucleus speckle. Nucleus, nuclear pore complex. Nucleus envelope. Cytoplasm. Cytoplasm, Stress granule. Note=Localized predominantly in the nucleoplasm and at both the nucleoplasmic and cytoplasmic faces of the nuclear pore complex. Shuttles between the nucleus and the cytoplasm. Travels to the cytoplasm as part of the exon junction complex (EJC) bound to mRNA. The association with the TREX complex seems to occur in regions surrounding nuclear speckles known as perispeckles (PubMed:23826332). Nucleus; nuclear rim (PubMed:25662211)

#### Tissue Location

Expressed ubiquitously.

#### NXF1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### NXF1 Rabbit mAb - Images



