

p21 Rabbit mAb

Catalog # AP75848

Specification

p21 Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB <u>P38936</u> Human, Mouse Rabbit Monoclonal Antibody 18119

p21 Rabbit mAb - Additional Information

Gene ID 1026

Other Names CDKN1A

Dilution WB~~1/500-1/1000

Format Liquid

p21 Rabbit mAb - Protein Information

Name CDKN1A (HGNC:1784)

Function

Plays an important role in controlling cell cycle progression and DNA damage-induced G2 arrest (PubMed:9106657). Involved in p53/TP53 mediated inhibition of cellular proliferation in response to DNA damage. Also involved in p53-independent DNA damage-induced G2 arrest mediated by CREB3L1 in astrocytes and osteoblasts (By similarity). Binds to and inhibits cyclin-dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D-CDK4 complex. Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (PubMed:11595739). Negatively regulates the CDK4- and CDK6-driven phosphorylation of RB1 in keratinocytes, thereby resulting in the release of E2F1 and subsequent transcription of E2F1-driven G1/S phase promoting genes (By similarity).

Cellular Location Cytoplasm. Nucleus

Tissue Location



Expressed in all adult tissues, with 5-fold lower levels observed in the brain

p21 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

p21 Rabbit mAb - Images

