

**PERK Rabbit mAb**  
Catalog # AP75893**Specification****PERK Rabbit mAb - Product Information**

Application	WB
Primary Accession	<a href="#">Q9NZJ5</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	125216

**PERK Rabbit mAb - Additional Information**

Gene ID 9451

**Other Names**  
EIF2AK3**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**PERK Rabbit mAb - Protein Information****Name** EIF2AK3 {ECO:0000303|PubMed:10932183, ECO:0000312|HGNC:HGNC:3255}**Function**

Metabolic-stress sensing protein kinase that phosphorylates the alpha subunit of eukaryotic translation initiation factor 2 (EIF2S1/eIF-2-alpha) in response to various stress, such as unfolded protein response (UPR) (PubMed: [10026192](http://www.uniprot.org/citations/10026192) target="\_blank">10026192</a>, PubMed: [10677345](http://www.uniprot.org/citations/10677345) target="\_blank">10677345</a>, PubMed: [11907036](http://www.uniprot.org/citations/11907036) target="\_blank">11907036</a>, PubMed: [12086964](http://www.uniprot.org/citations/12086964) target="\_blank">12086964</a>, PubMed: [25925385](http://www.uniprot.org/citations/25925385) target="\_blank">25925385</a>, PubMed: [31023583](http://www.uniprot.org/citations/31023583) target="\_blank">31023583</a>). Key effector of the integrated stress response (ISR) to unfolded proteins: EIF2AK3/PERK specifically recognizes and binds misfolded proteins, leading to its activation and EIF2S1/eIF-2-alpha phosphorylation (PubMed: [10677345](http://www.uniprot.org/citations/10677345) target="\_blank">10677345</a>, PubMed: [27917829](http://www.uniprot.org/citations/27917829) target="\_blank">27917829</a>, PubMed: [31023583](http://www.uniprot.org/citations/31023583) target="\_blank">31023583</a>). EIF2S1/eIF-2-alpha phosphorylation in response to stress converts EIF2S1/eIF-2-alpha in a global protein synthesis inhibitor, leading to a global attenuation of cap-dependent translation, while concomitantly initiating the preferential translation of ISR-specific mRNAs, such as the transcriptional activators ATF4 and QRICH1, and hence allowing ATF4- and QRICH1-mediated

reprogramming (PubMed:<a href="http://www.uniprot.org/citations/10026192" target="\_blank">10026192</a>, PubMed:<a href="http://www.uniprot.org/citations/10677345" target="\_blank">10677345</a>, PubMed:<a href="http://www.uniprot.org/citations/31023583" target="\_blank">31023583</a>, PubMed:<a href="http://www.uniprot.org/citations/33384352" target="\_blank">33384352</a>). The EIF2AK3/PERK- mediated unfolded protein response increases mitochondrial oxidative phosphorylation by promoting ATF4-mediated expression of COX7A2L/SCAF1, thereby increasing formation of respiratory chain supercomplexes (PubMed:<a href="http://www.uniprot.org/citations/31023583" target="\_blank">31023583</a>). In contrast to most subcellular compartments, mitochondria are protected from the EIF2AK3/PERK-mediated unfolded protein response due to EIF2AK3/PERK inhibition by ATAD3A at mitochondria-endoplasmic reticulum contact sites (PubMed:<a href="http://www.uniprot.org/citations/39116259" target="\_blank">39116259</a>). In addition to EIF2S1/eIF-2-alpha, also phosphorylates NFE2L2/NRF2 in response to stress, promoting release of NFE2L2/NRF2 from the BCR(KEAP1) complex, leading to nuclear accumulation and activation of NFE2L2/NRF2 (By similarity). Serves as a critical effector of unfolded protein response (UPR)-induced G1 growth arrest due to the loss of cyclin-D1 (CCND1) (By similarity). Involved in control of mitochondrial morphology and function (By similarity).

#### Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q9Z2B5}; Single-pass type I membrane protein. Note=Localizes to the Localizes to endoplasmic reticulum membrane (By similarity). Also present at mitochondria-endoplasmic reticulum contact sites; where it interacts with ATAD3A (PubMed:39116259). {ECO:0000250|UniProtKB:Q9Z2B5, ECO:0000269|PubMed:39116259}

#### Tissue Location

Ubiquitous. A high level expression is seen in secretory tissues.

#### PERK Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### PERK Rabbit mAb - Images



