

**PGK1 Rabbit mAb**  
Catalog # AP75900**Specification****PGK1 Rabbit mAb - Product Information**

Application	WB, IF
Primary Accession	<a href="#">P00558</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	44615

**PGK1 Rabbit mAb - Additional Information**

Gene ID 5230

**Other Names**  
PGK1**Dilution**  
WB~~1/500-1/1000  
IF~~1/50-1/200**Format**  
Liquid**PGK1 Rabbit mAb - Protein Information****Name** PGK1**Synonyms** PGKA**Function**

Catalyzes one of the two ATP producing reactions in the glycolytic pathway via the reversible conversion of 1,3- diphosphoglycerate to 3-phosphoglycerate (PubMed:<a href="http://www.uniprot.org/citations/30323285" target="\_blank">30323285</a>, PubMed:<a href="http://www.uniprot.org/citations/7391028" target="\_blank">7391028</a>). Both L- and D-forms of purine and pyrimidine nucleotides can be used as substrates, but the activity is much lower on pyrimidines (PubMed:<a href="http://www.uniprot.org/citations/18463139" target="\_blank">18463139</a>). In addition to its role as a glycolytic enzyme, it seems that PGK1 acts as a polymerase alpha cofactor protein (primer recognition protein) (PubMed:<a href="http://www.uniprot.org/citations/2324090" target="\_blank">2324090</a>). Acts as a protein kinase when localized to the mitochondrion where it phosphorylates pyruvate dehydrogenase kinase PDK1 to inhibit pyruvate dehydrogenase complex activity and suppress the formation of acetyl- coenzyme A from pyruvate, and consequently inhibit oxidative phosphorylation and promote glycolysis (PubMed:<a href="http://www.uniprot.org/citations/26942675" target="\_blank">26942675</a>, PubMed:<a href="http://www.uniprot.org/citations/36849569" target="\_blank">36849569</a>). May play a

role in sperm motility (PubMed:<a href="http://www.uniprot.org/citations/26677959" target="\_blank">26677959</a>).

#### Cellular Location

Cytoplasm, cytosol. Mitochondrion matrix. Note=Hypoxic conditions promote mitochondrial targeting (PubMed:26942675). Targeted to the mitochondrion following phosphorylation by MAPK1/ERK2, cis-trans isomerization by PIN1, and binding to mitochondrial circRNA mcPGK1 (PubMed:36849569).

#### Tissue Location

Mainly expressed in spermatogonia. Localized on the principle piece in the sperm (at protein level). Expression significantly decreased in the testis of elderly men

### PGK1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### PGK1 Rabbit mAb - Images



