

**Ran Rabbit mAb**  
Catalog # AP76001**Specification**

---

**Ran Rabbit mAb - Product Information**

Application	<b>WB, IF, IP</b>
Primary Accession	<a href="#">P62826</a>
Reactivity	<b>Human, Monkey</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>24423</b>

**Ran Rabbit mAb - Additional Information**

Gene ID 5901

**Other Names**

RAN

**Dilution**

WB~~1/500-1/1000

IF~~1/50-1/200

IP~~1/20

**Format**

Liquid

**Ran Rabbit mAb - Protein Information**

Name RAN

**Synonyms** ARA24 {ECO:0000303|PubMed:10400640}**Function**

GTPase involved in nucleocytoplasmic transport, participating both to the import and the export from the nucleus of proteins and RNAs (PubMed:[10400640](http://www.uniprot.org/citations/10400640), PubMed:[17209048](http://www.uniprot.org/citations/17209048), PubMed:[26272610](http://www.uniprot.org/citations/26272610), PubMed:[27306458](http://www.uniprot.org/citations/27306458), PubMed:[8276887](http://www.uniprot.org/citations/8276887), PubMed:[8636225](http://www.uniprot.org/citations/8636225), PubMed:[8692944](http://www.uniprot.org/citations/8692944), PubMed:[8896452](http://www.uniprot.org/citations/8896452), PubMed:[9351834](http://www.uniprot.org/citations/9351834), PubMed:[9428644](http://www.uniprot.org/citations/9428644), PubMed:[9822603](http://www.uniprot.org/citations/9822603)). Switches between a cytoplasmic GDP- and a nuclear GTP-bound state by nucleotide exchange and GTP

hydrolysis (PubMed:<a href="http://www.uniprot.org/citations/11336674" target="\_blank">11336674</a>, PubMed:<a href="http://www.uniprot.org/citations/26272610" target="\_blank">26272610</a>, PubMed:<a href="http://www.uniprot.org/citations/29040603" target="\_blank">29040603</a>, PubMed:<a href="http://www.uniprot.org/citations/7819259" target="\_blank">7819259</a>, PubMed:<a href="http://www.uniprot.org/citations/8636225" target="\_blank">8636225</a>, PubMed:<a href="http://www.uniprot.org/citations/8692944" target="\_blank">8692944</a>, PubMed:<a href="http://www.uniprot.org/citations/8896452" target="\_blank">8896452</a>, PubMed:<a href="http://www.uniprot.org/citations/9351834" target="\_blank">9351834</a>, PubMed:<a href="http://www.uniprot.org/citations/9428644" target="\_blank">9428644</a>, PubMed:<a href="http://www.uniprot.org/citations/9822603" target="\_blank">9822603</a>). Nuclear import receptors such as importin beta bind their substrates only in the absence of GTP-bound RAN and release them upon direct interaction with GTP-bound RAN, while export receptors behave in the opposite way. Thereby, RAN controls cargo loading and release by transport receptors in the proper compartment and ensures the directionality of the transport (PubMed:<a href="http://www.uniprot.org/citations/8896452" target="\_blank">8896452</a>, PubMed:<a href="http://www.uniprot.org/citations/9351834" target="\_blank">9351834</a>, PubMed:<a href="http://www.uniprot.org/citations/9428644" target="\_blank">9428644</a>). Interaction with RANBP1 induces a conformation change in the complex formed by XPO1 and RAN that triggers the release of the nuclear export signal of cargo proteins (PubMed:<a href="http://www.uniprot.org/citations/20485264" target="\_blank">20485264</a>). RAN (GTP-bound form) triggers microtubule assembly at mitotic chromosomes and is required for normal mitotic spindle assembly and chromosome segregation (PubMed:<a href="http://www.uniprot.org/citations/10408446" target="\_blank">10408446</a>, PubMed:<a href="http://www.uniprot.org/citations/29040603" target="\_blank">29040603</a>). Required for normal progress through mitosis (PubMed:<a href="http://www.uniprot.org/citations/12194828" target="\_blank">12194828</a>, PubMed:<a href="http://www.uniprot.org/citations/29040603" target="\_blank">29040603</a>, PubMed:<a href="http://www.uniprot.org/citations/8421051" target="\_blank">8421051</a>). The complex with BIRC5/survivin plays a role in mitotic spindle formation by serving as a physical scaffold to help deliver the RAN effector molecule TPX2 to microtubules (PubMed:<a href="http://www.uniprot.org/citations/18591255" target="\_blank">18591255</a>). Acts as a negative regulator of the kinase activity of VRK1 and VRK2 (PubMed:<a href="http://www.uniprot.org/citations/18617507" target="\_blank">18617507</a>). Enhances AR- mediated transactivation. Transactivation decreases as the poly-Gln length within AR increases (PubMed:<a href="http://www.uniprot.org/citations/10400640" target="\_blank">10400640</a>).

### Cellular Location

Nucleus. Nucleus envelope. Cytoplasm, cytosol Cytoplasm. Melanosome Note=Predominantly nuclear during interphase (PubMed:10679025, PubMed:12194828, PubMed:8421051). Becomes dispersed throughout the cytoplasm during mitosis (PubMed:12194828, PubMed:8421051). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065).

### Tissue Location

Expressed in a variety of tissues.

### Ran Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Ran Rabbit mAb - Images



