

# Complex III Subunit 5 Rabbit mAb

Catalog # AP76024

### Specification

## Complex III Subunit 5 Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC, IF <u>P47985</u> Human, Mouse, Rat Rabbit Monoclonal Antibody 29668

### Complex III Subunit 5 Rabbit mAb - Additional Information

Gene ID 7386

Other Names UQCRFS1

Dilution WB~~1/500-1/1000 IHC~~1/50-1/100 IF~~1/50-1/200

Format Liquid

## **Complex III Subunit 5 Rabbit mAb - Protein Information**

Name UQCRFS1 (HGNC:12587)

#### Function

[Cytochrome b-c1 complex subunit Rieske, mitochondrial]: Component of the ubiquinol-cytochrome c oxidoreductase, a multisubunit transmembrane complex that is part of the mitochondrial electron transport chain which drives oxidative phosphorylation (PubMed:<a href="http://www.uniprot.org/citations/31883641" target=" blank">31883641</a>). The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol- cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. The cytochrome b- c1 complex catalyzes electron transfer from ubiquinol to cytochrome c, linking this redox reaction to translocation of protons across the mitochondrial inner membrane, with protons being carried across the membrane as hydrogens on the quinol. In the process called Q cycle, 2 protons are consumed from the matrix, 4 protons are released into the intermembrane space and 2 electrons are passed to cytochrome c. The Rieske protein is a catalytic core subunit containing a [2Fe-2S] iron- sulfur cluster. It cycles between 2 conformational states during catalysis to transfer electrons from the quinol bound in the Q(0) site in cytochrome b to cytochrome c1 (By similarity).



Incorporation of UQCRFS1 is the penultimate step in complex III assembly (PubMed:<a href="http://www.uniprot.org/citations/28673544" target="\_blank">28673544</a>).

**Cellular Location** Mitochondrion inner membrane; Single-pass membrane protein {ECO:0000250|UniProtKB:Q5ZLR5}

## **Complex III Subunit 5 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### Complex III Subunit 5 Rabbit mAb - Images





