

# S100A4 Rabbit mAb

Catalog # AP76046

### Specification

# S100A4 Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC, IF <u>P26447</u> Human Rabbit Monoclonal Antibody 11729

### S100A4 Rabbit mAb - Additional Information

Gene ID 6275

Other Names S100A4

**Dilution** WB~~1/500-1/1000 IHC~~1/50-1/100 IF~~1/50-1/200

Format Liquid

# S100A4 Rabbit mAb - Protein Information

Name S100A4

Synonyms CAPL, MTS1

Function

Calcium-binding protein that plays a role in various cellular processes including motility, angiogenesis, cell differentiation, apoptosis, and autophagy (PubMed:<a

href="http://www.uniprot.org/citations/16707441" target="\_blank">16707441</a>, PubMed:<a href="http://www.uniprot.org/citations/23752197" target="\_blank">23752197</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">30713770</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">30713770</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">16707441</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">30713770</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">30713770</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">16707441</a> IIA/MYH9 (PubMed:<a href="http://www.uniprot.org/citations/16707441"

target="\_blank">16707441</a>). Mechanistically, promotes filament depolymerization and increases the amount of soluble myosin-IIA, resulting in the formation of stable protrusions facilitating chemotaxis (By similarity). Modulates also the pro-apoptotic function of TP53 by binding to its C-terminal transactivation domain within the nucleus and reducing its protein levels (PubMed:<a href="http://www.uniprot.org/citations/23752197" target="\_blank">23752197</a>). Within the extracellular space, stimulates cytokine production including granulocyte colony-stimulating factor and CCL24 from T-lymphocytes (By similarity). In addition, stimulates



T-lymphocyte chemotaxis by acting as a chemoattractant complex with PGLYRP1 that promotes lymphocyte migration via CCR5 and CXCR3 receptors (PubMed:<a href="http://www.uniprot.org/citations/26654597" target="\_blank">26654597</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">30713770</a>).

**Cellular Location** Secreted. Nucleus Cytoplasm {ECO:0000250|UniProtKB:P07091}

**Tissue Location** Ubiquitously expressed.

### S100A4 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- S100A4 Rabbit mAb Images



